

BRICK POLYTOPES OF SPHERICAL SUBWORD COMPLEXES: A NEW APPROACH TO GENERALIZED ASSOCIAHEDRA

VINCENT PILAUD AND CHRISTIAN STUMP

ABSTRACT. We generalize the brick polytope of V. Pilaud and F. Santos to spherical subword complexes for finite Coxeter groups. This construction provides polytopal realizations for a certain class of subword complexes containing all cluster complexes of finite types. For the latter, the brick polytopes turn out to coincide with the known realizations of generalized associahedra, thus opening new perspectives on these constructions. This new approach yields in particular the vertex description of generalized associahedra, and a Minkowski sum decomposition into Coxeter matroid polytopes.

KEYWORDS. Coxeter–Catalan combinatorics, subword complexes, cluster complexes, generalized associahedra, Cambrian lattices, Cambrian fans

CONTENTS

Introduction	2
1. Main results	3
1.1. Main definitions	3
1.2. Realizing words	3
1.3. Further properties	4
1.4. Generalized associahedra	4
2. Background on Coxeter groups and subword complexes	5
2.1. Finite Coxeter groups	5
2.2. Subword complexes	8
3. Root configurations	12
4. Generalized brick polytopes	15
5. Further combinatorial and geometric properties	18
5.1. A surjective map	18
5.2. A normal fan description	20
5.3. Increasing flips and quotients of the weak order	21
5.4. Fibers of κ and the weak order	22
5.5. Canonical spanning trees of the increasing flip graph	24
5.6. Minkowski sum decompositions into Coxeter matroid polytopes	26
6. Cluster complexes and generalized associahedra revisited	27
6.1. Realizing cluster complexes	28
6.2. Normal vectors and facets	29
6.3. Five Coxeter-Catalan families	34
6.4. Cambrian lattices and fans	41
7. Conclusion	42
Acknowledgements	44
References	44

Vincent Pilaud was partially supported by grants MTM2008-04699-C03-02 and MTM2011-22792 of the spanish MICINN, by European Research Project ExploreMaps (ERC StG 208471), and by a postdoctoral grant of the Fields Institute of Toronto.

Christian Stump was partially supported by a CRM-ISM fellowship at UQAM, Montreal, and by the DFG via the Research Group “Methods for Discrete Structures”.

INTRODUCTION

In [PP12], the first author and M. Pocchiola studied pseudoline arrangements on sorting networks as combinatorial models for triangulations of convex polygons, pseudotriangulations [RSS08], and multitriangulations [PS09]. Motivated by the question of the polytopality of the multiassociahedron, the first author and F. Santos [PS12] defined the brick polytope of a sorting network. This construction provides polytopal realizations of the simplicial complexes associated to a certain class of sorting networks. In particular, it turns out that all type A associahedra of C. Hohlweg and C. Lange [HL07] appear as brick polytopes for well-chosen sorting networks.

Independently in [Stu11], the second author connected multitriangulations and subword complexes of type A , which can be visually interpreted with sorting networks (see also a generalization developed in [SS12]). Subword complexes were introduced by A. Knutson and E. Miller in the context of Gröbner geometry of Schubert varieties [KM05]. The definition was then extended to all Coxeter groups in [KM04]. They proved that these simplicial complexes are either topological balls or spheres, and raised the question of realizing spherical subword complexes as boundary complexes of convex polytopes.

Recently, C. Ceballos, J.-P. Labbé, and the second author [CLS11] extended the subword complex interpretations of [Stu11] and [PP12] for triangulations and multitriangulations of convex polygons to general finite types. They proved in particular that all finite type cluster complexes of S. Fomin and A. Zelevinsky [FZ03] are isomorphic to certain well-chosen subword complexes. The first polytopal realizations of cluster complexes of finite types were obtained by F. Chapoton, S. Fomin and A. Zelevinsky [CFZ02]. Later, C. Hohlweg, C. Lange and H. Thomas [HLT11] extended the ideas of [HL07] to construct different realizations based on the Cambrian fan studied by N. Reading and D. Speyer [Rea04, Rea06, Rea07b, RS09]. In his recent preprint [Ste11], S. Stella generalizes the approach of [CFZ02] and shows that the resulting realizations of the associahedra coincide with that of [HLT11].

In this paper, we define the brick polytope of any subword complex for a finite Coxeter group. As in type A , the brick polytope does not realize its subword complex in general. We study the class $\mathfrak{R}$ of subword complexes which are realized by their brick polytope. They are simply characterized in terms of their root configuration. The class $\mathfrak{R}$ contains in particular the finite type cluster complexes. For these, the brick polytopes turn out to be (translates of) the realizations of the generalized associahedra described in [CFZ02, HLT11, Ste11]. We therefore obtain the vertex description of these realizations and decompose them into Minkowski sums of Coxeter matroid polytopes as defined by A. Borovik, I. Gelfand, and N. White [BGW97]. We moreover revisit the combinatorics of Cambrian fans, and provide a detailed picture of the bijective connections between clusters, sortable elements, noncrossing partitions and vertices of the brick polytope. Furthermore, we extend relevant combinatorial properties of these realizations of the cluster complex to any subword complex of the class $\mathfrak{R}$. We relate the normal fan of the brick polytope to the Coxeter fan, and the graph of the brick polytope to a quotient of the weak order.

Along the paper, we rephrase our main definitions in the classical situation of type A to show that they match the definitions in [PS12], more details can be found therein. The reader may as well skip these parts for a more compact presentation.

1. MAIN RESULTS

In this section, we summarize the main results of the paper for the convenience of the reader. Precise definitions and detailed proofs appear in further sections.

1.1. Main definitions. Consider a finite Coxeter system (W, S) , with simple roots $\Delta := \{\alpha_s \mid s \in S\}$, and fundamental weights $\nabla := \{\omega_s \mid s \in S\}$. The group W acts on the vector space V with basis Δ . For a word $Q := q_1 \cdots q_m$ in S^* and an element $\rho \in W$, A. Knutson and E. Miller [KM04] define the subword complex $\mathcal{S}(Q, \rho)$ to be the simplicial complex of subwords of Q whose complement contains a reduced expression of ρ . Without loss of generality, we assume that $\rho = w_\circ = \delta(Q)$, where $w_\circ$ denotes the longest element of W and $\delta(Q)$ denotes the Demazure product of Q . The subword complex $\mathcal{S}(Q) := \mathcal{S}(Q, w_\circ)$ is then a topological sphere [KM04].

After recalling some background on Coxeter groups and subword complexes in Section 2, we study in Sections 3 and 4 the root and the weight function on facets in a subword complex. Associate to any facet I of the subword complex $\mathcal{S}(Q)$ a root function $r(I, \cdot) : [m] \rightarrow W(\Delta)$ and a weight function $w(I, \cdot) : [m] \rightarrow W(\nabla)$ defined by

$$r(I, k) := \sigma_I^{[1, k-1]}(\alpha_{q_k}) \quad \text{and} \quad w(I, k) := \sigma_I^{[1, k-1]}(\omega_{q_k}),$$

where $\sigma_I^{[x, y]}$ denotes the product of the reflections q_z for $z \in [x, y] \setminus I$ in this order. The root function locally encodes the flip property in the subword complex: each facet adjacent to I in $\mathcal{S}(Q)$ is obtained by exchanging an element $i \in I$ with the unique element $j \notin I$ such that $r(I, j) \in \{\pm r(I, i)\}$.

We use the root function to define the *root configuration* of I as the multiset

$$R(I) := \{\{r(I, i) \mid i \in I\}\}.$$

On the other hand, we use the weight function to define the *brick vector* of I as

$$B(I) := \sum_{k \in [m]} w(I, k),$$

and the *brick polytope* of Q as the convex hull of the brick vectors of all facets of the subword complex $\mathcal{S}(Q)$,

$$\mathcal{B}(Q) := \text{conv} \{B(I) \mid I \text{ facet of } \mathcal{S}(Q)\}.$$

In the particular situation of type A Coxeter groups, these definitions match that of [PS12] for brick polytopes of sorting networks.

1.2. Realizing words. In this paper, we focus on *realizing* words, *i.e.* words Q for which the root configuration $R(I)$ of a facet I of $\mathcal{S}(Q)$ (or equivalently of all facets of $\mathcal{S}(Q)$) forms a linear basis for the vector space V . For these words, we obtain the main result of this paper in Section 4.

Theorem 1.1. *If Q is a realizing word, then the polar of the brick polytope $\mathcal{B}(Q)$ realizes the subword complex $\mathcal{S}(Q)$.*

This theorem relies on elementary lemmas which relate the root configuration of a facet I of $\mathcal{S}(Q)$ with the edges of $\mathcal{B}(Q)$ incident to $B(I)$, as follows.

Proposition 1.2. *The cone of the brick polytope $\mathcal{B}(Q)$ at the brick vector $B(I)$ of a facet I of $\mathcal{S}(Q)$ is generated by the negative of the root configuration $R(I)$, *i.e.**

$$\text{cone} \{B(J) - B(I) \mid J \text{ facet of } \mathcal{S}(Q)\} = \text{cone} \{-r(I, i) \mid i \in I\}.$$

1.3. Further properties. Further combinatorial and geometric properties of our construction are developed in Section 5. We first define a map κ from the Coxeter group W to the facets of the subword complex $\mathcal{S}(Q)$ which associates to an element $w \in W$ the unique facet $\kappa(w)$ of $\mathcal{S}(Q)$ whose root configuration $R(\kappa(w))$ is a subset of $w(\Phi^+)$. In other words, $\kappa(w)$ is the facet of $\mathcal{S}(Q)$ whose brick vector $B(\kappa(w))$ maximizes the functional $x \mapsto \langle w(q)|x \rangle$, where q is any point in the interior of the fundamental chamber $\mathcal{C}$ of W . The map κ thus enables us to describe the normal fan of the brick polytope. It is obtained by gluing the chambers of the Coxeter fan according to the fibers of κ as follows.

Proposition 1.3. *The normal cone of $B(I)$ in $\mathcal{B}(Q)$ is the union of the chambers $w(\mathcal{C})$ of the Coxeter fan of W given by the elements $w \in W$ with $\kappa(w) = I$.*

We then relate the graph of the brick polytope $\mathcal{B}(Q)$ to the weak order on W . Let I and J be two adjacent facets of $\mathcal{S}(Q)$ with $I \setminus i = J \setminus j$. We say that the flip from I to J is *increasing* if $i < j$. We call *increasing flip order* the transitive closure of the graph of increasing flips.

Proposition 1.4. *A facet I is covered by a facet J in increasing flip order if and only if there exist $w_I \in \kappa^{-1}(I)$ and $w_J \in \kappa^{-1}(J)$ such that w_I is covered by w_J in weak order.*

Based on computational evidence, we conjecture that the increasing flip order defines a lattice structure on the facets of the subword complex $\mathcal{S}(Q)$. In relation to this question, we study some properties of the fibers of the map κ with respect to the weak order: we show that the fibers are closed by intervals and we characterize the fibers which admit a meet or a join. Finally, we describe a canonical oriented spanning tree of the increasing flip graph.

To close our study of brick polytopes of realizing words, we provide a Minkowski sum decomposition of these realizations into Coxeter matroid polytopes [BGW97].

Proposition 1.5. *The brick polytope $\mathcal{B}(Q)$ is the Minkowski sum of the polytopes*

$$\mathcal{B}(Q, k) := \text{conv} \{w(I, k) \mid I \text{ facet of } \mathcal{S}(Q)\}$$

over all positions $k \in [m]$. Moreover, each summand $\mathcal{B}(Q, k)$ is a Coxeter matroid polytope as defined in [BGW97].

1.4. Generalized associahedra. Our last section is devoted to the main motivation and example of brick polytopes. Fix a Coxeter element c of W , and a reduced expression c of c . We denote by $w_\circ(c)$ the c -sorting word of $w_\circ$. As recently proved by [CLS11], the subword complex $\mathcal{S}(cw_\circ(c))$ is then isomorphic to the c -cluster complex of W as defined by N. Reading in [Rea07a]. Since the word $cw_\circ(c)$ is realizing, we can apply our brick polytope construction to obtain polytopal realizations of the cluster complex.

Theorem 1.6. *For any reduced expression c of any Coxeter element c of W , the polar of the brick polytope $\mathcal{B}(cw_\circ(c))$ realizes the subword complex $\mathcal{S}(cw_\circ(c))$, and thus the c -cluster complex of type W .*

We prove that brick polytopes are in fact translates of the known realizations of the generalized associahedra [CFZ02, HLT11, Ste11]. We thus obtain the vertex description of these realizations, and a Minkowski sum decomposition into Coxeter matroid polytopes. Furthermore, we obtain independent elementary proofs of the

following two results, which are the heart of the construction of [HLT11]. We say that an element $w \in W$ is a *singleton* if $\kappa^{-1}(\kappa(w)) = \{w\}$.

Proposition 1.7. *Up to translation by a vector Ω , the brick polytope $\mathcal{B}(\text{cw}_\circ(c))$ is obtained from the balanced W -permutahedron $\mathcal{P}(W)$ removing all facets which do not intersect the set $\{w(\sum_s \omega_s) \mid w \in W \text{ singleton}\}$.*

Proposition 1.8. *The following properties are equivalent for an element $w \in W$:*

- (i) *The element w is a singleton.*
- (ii) *The root configuration $R(\kappa(w)) := \{r(\kappa(w), i) \mid i \in \kappa(w)\}$ equals $w(\Delta)$.*
- (iii) *The weight configuration $W(\kappa(w)) := \{w(\kappa(w), i) \mid i \in \kappa(w)\}$ equals $w(\nabla)$.*
- (iv) *The vertices $B(\kappa(w))$ of the brick polytope $\mathcal{B}(\text{cw}_\circ(c))$ and $w(q)$ of the balanced W -permutahedron $\mathcal{P}(W)$ coincide up to Ω , i.e. $B(\kappa(w)) = \Omega + w(q)$.*
- (v) *There exist reduced expressions w of w and c of c such that w is a prefix of $w_\circ(c)$.*
- (vi) *The complement of $\kappa(w)$ in $\text{cw}_\circ(c)$ is $w_\circ(c)$.*

Finally, we revisit the connection between these polytopal realizations of the cluster complex and the Cambrian lattices and fans defined by N. Reading and D. Speyer in [Rea06, Rea07b, RS09]. We first describe in detail the bijective connections between the facets of the subword complex $\mathcal{S}(\text{cw}_\circ(c))$ and different Coxeter-Catalan families studied in [Rea07b]: c -clusters, c -sortable elements, and c -noncrossing partitions and subspaces. In particular, combining the bijection of [CLS11] from the subword complex $\mathcal{S}(\text{cw}_\circ(c))$ to the c -cluster complex with the bijection of [Rea07b] from c -sortable elements in W to the cluster complex, we describe a bijection so_c from facets of $\mathcal{S}(\text{cw}_\circ(c))$ to c -sortable elements. We then prove that the set $\mathcal{C}_c(w)$ defined in [RS09] coincides with the root configuration $R(\text{so}_c^{-1}(w))$, and that the map so_c^{-1} is the restriction of κ to c -sortable elements. This observation yields the following results.

Proposition 1.9. *If $v, w \in W$ and v is c -sortable, then $\kappa(w) = \kappa(v)$ if and only if v is the maximal c -sortable element below w in weak order.*

Corollary 1.10. *For the word $\text{cw}_\circ(c)$, the bijection so_c is an isomorphism between the Cambrian lattice on c -sortable elements in W and the increasing flip order on the facets of $\mathcal{S}(\text{cw}_\circ(c))$.*

Corollary 1.11. *The normal fan of the brick polytope $\mathcal{B}(\text{cw}_\circ(c))$ coincides with the c -Cambrian fan.*

Although Corollary 1.10 follows from [IS10, IT09] for simply laced types and Corollary 1.11 could be deduced from the connection between the brick polytope $\mathcal{B}(\text{cw}_\circ(c))$ and the c -associahedron of [HLT11], the subword complex approach provides new independent proofs.

2. BACKGROUND ON COXETER GROUPS AND SUBWORD COMPLEXES

2.1. Finite Coxeter groups. We recall here classical notions on finite Coxeter groups. See [Hum90, Hum78] for more details.

2.1.1. Coxeter systems. Let $(V, \langle \cdot | \cdot \rangle)$ be an n -dimensional euclidean vector space. For any vector $v \in V \setminus 0$, we denote by s_v the reflection interchanging v and $-v$ while fixing the orthogonal hyperplane pointwise. Remember that $ws_v = s_{w(v)}w$ for any non-zero vector $v \in V$ and any orthogonal transformation w .

We consider a *finite Coxeter group* W acting on V , that is, a finite group generated by reflections. The set of *reflections* in W is denoted by R . The *Coxeter arrangement* of W is the collection of all reflecting hyperplanes. Its complement in V is a union of open polyhedral cones. Their closures are called *chambers*. The *Coxeter fan* is the polyhedral fan formed by the chambers together with all their faces. This fan is *complete* (its cones cover V) and *simplicial* (all cones are simplicial), and we can assume without loss of generality that it is *essential* (the intersection of all chambers is reduced to the origin). We fix an arbitrary chamber $\mathcal{C}$ which we call the *fundamental chamber*. The *simple reflections* of W are the n reflections orthogonal to the facet defining hyperplanes of $\mathcal{C}$. The set $S \subseteq R$ of simple reflections generates W . In particular, $R = \{ws w^{-1} \mid w \in W, s \in S\}$. The pair (W, S) forms a *Coxeter system*.

For inductive procedures, set $W_{\langle s \rangle}$ for $s \in S$ to be the *parabolic subgroup* of W generated by $S \setminus s$. It is a Coxeter group with Coxeter system $(W_{\langle s \rangle}, S \setminus s)$.

2.1.2. Roots and weights. For simple reflections $s, t \in S$, denote by m_{st} the order of the product $st \in W$. We fix a *generalized Cartan matrix* for (W, S) , i.e. a matrix $[a_{st}]_{s, t \in S}$ such that $a_{ss} = 2$, $a_{st} \leq 0$, $a_{st}a_{ts} = 4 \cos^2(\frac{\pi}{m_{st}})$ and $a_{st} = 0 \Leftrightarrow a_{ts} = 0$ for all $s, t \in S$. We can associate to each simple reflection s a *simple root* $\alpha_s \in V$, orthogonal to the reflecting hyperplane of s and pointing toward the half-space containing $\mathcal{C}$, in such a way that $s(\alpha_t) = \alpha_t - a_{st}\alpha_s$ for all $s, t \in S$. The set of all simple roots is denoted by $\Delta := \{\alpha_s \mid s \in S\}$. The orbit $\Phi := \{w(\alpha_s) \mid w \in W, s \in S\}$ of Δ under W is a *root system* for W . It is invariant under the action of W and contains precisely two opposite roots orthogonal to each reflecting hyperplane of W .

The set Δ of simple roots forms a linear basis of V (since we assumed W to act essentially on V). The root system Φ is the disjoint union of the *positive roots* $\Phi^+ := \Phi \cap \mathbb{R}_{\geq 0}[\Delta]$ (non-negative linear combinations of the simple roots) and the *negative roots* $\Phi^- := -\Phi^+$. In other words, the positive roots are the roots whose scalar product with any vector of the interior of the fundamental chamber $\mathcal{C}$ is positive, and the simple roots form the basis of the cone generated by Φ^+ . Each reflection hyperplane is orthogonal to one positive and one negative root. For a reflection $s \in R$, we set α_s to be the unique positive root orthogonal to the reflection hyperplane of s , i.e. such that $s = s_{\alpha_s}$. For $\alpha \in \Delta$, we also denote by

$$\Phi_{\langle \alpha \rangle} := \Phi \cap \text{vect}(\Delta \setminus \alpha)$$

the *parabolic subroot system* of Φ generated by $\Delta \setminus \{\alpha\}$ with corresponding parabolic subgroup $W_{\langle s_\alpha \rangle}$.

The set of *fundamental weights* $\nabla := \{\omega_s \mid s \in S\}$ of the Coxeter system (W, S) is the dual basis to the renormalized root basis $\{2\alpha_s / \langle \alpha_s | \alpha_s \rangle \mid s \in S\}$. In other words, the weights are defined by $\langle \alpha_s | \omega_t \rangle = \delta_{s=t} \langle \alpha_s | \alpha_s \rangle / 2$ for all $s, t \in S$. Thus, the Coxeter group W acts on the weight basis by $s(\omega_t) = \omega_t - \delta_{s=t} \alpha_s$. Note that the transposed Cartan matrix transforms the weight basis into the root basis, i.e. $\alpha_s = \sum_{t \in S} a_{ts} \omega_t$ for any $s \in S$. Geometrically, the weight ω_s gives the direction of the ray of the fundamental chamber $\mathcal{C}$ not contained in the reflecting hyperplane of s .

2.1.3. Coxeter permutahedra. The *W-permutahedron* $\mathcal{P}_q(W)$ is the convex hull of the orbit under W of a point q in the interior of the fundamental chamber $\mathcal{C}$ of W . Its normal fan is the *Coxeter fan* of W . The facet of $\mathcal{P}_q(W)$ orthogonal to $w(\omega_s)$ is defined by the inequality $\langle w(\omega_s) | x \rangle \leq \langle \omega_s | q \rangle$ and is supported by the

hyperplane $w(q + \text{vect}(\Delta \setminus \alpha_s))$. If $q = \sum_{s \in S} \omega_s$, then $\mathcal{P}_q(W)$ is (a translate of) the Minkowski sum of all positive roots (each considered as a one-dimensional polytope). We call it the *balanced W -permutahedron*, and denote it by $\mathcal{P}(W)$.

2.1.4. Words on S . The *length* $\ell(w)$ of an element $w \in W$ is the length of the smallest expression of w as a product of the generators in S . An expression $w = w_1 \cdots w_p$ with $w_1, \dots, w_p \in S$ is called *reduced* if $p = \ell(w)$. Geometrically, the length of $w \in W$ is the number of positive roots sent to negative roots by w . Note that for any $w \in W$ and $s \in S$, we have $\ell(ws) = \ell(w) + 1$ if $w(\alpha_s) \in \Phi^+$ and $\ell(ws) = \ell(w) - 1$ if $w(\alpha_s) \in \Phi^-$. We denote by $w_\circ$ the unique *longest element* in W . It sends all positive roots to negatives.

The *(right) weak order* on W is the partial order $\leq$ defined by $u \leq w$ if there exists $v \in W$ such that $uv = w$ and $\ell(u) + \ell(v) = \ell(w)$. It defines a lattice structure on the elements of W with minimal element being the identity $e \in W$ and with maximal element being $w_\circ$. Its Hasse diagram is the graph of the W -permutahedron (oriented from e to $w_\circ$). The elements of W which cover a given element $w \in W$ in weak order are precisely the products ws for which $w(\alpha_s) \in \Phi^+$.

We denote by S^* the set of words on the alphabet S , and by ε the empty word. To avoid confusion, we denote with a square letter s the letter of the alphabet S corresponding to the single reflection $s \in S$. Similarly, we use a square letter like w to denote a word of S^* , and a normal letter like w to denote its corresponding group element in W . For example, we write $w := w_1 \cdots w_p$ meaning that the word $w \in S^*$ is formed by the letters $w_1, \dots, w_p$, while we write $w := w_1 \cdots w_p$ meaning that the element $w \in W$ is the product of the simple reflections $w_1, \dots, w_p$. The *Demazure product* on the Coxeter system (W, S) is the function $\delta : S^* \rightarrow W$ defined inductively by

$$\delta(\varepsilon) = e \quad \text{and} \quad \delta(Qs) = \begin{cases} ws & \text{if } \ell(ws) = \ell(w) + 1 \\ w & \text{if } \ell(ws) = \ell(w) - 1 \end{cases},$$

where $w = \delta(Q)$ denotes the Demazure product of Q . As its name suggests, $\delta(Q)$ corresponds to the product of the letters of Q in the Demazure algebra of (W, S) .

2.1.5. Crystallographic groups. The Coxeter group W is said to be *crystallographic* if it stabilizes a lattice of V . This can only happen if all entries of the Cartan matrix are integers. Reciprocally, if all entries of the Cartan matrix are integers, then the lattice generated by Δ is fixed by the Coxeter group W .

Example 2.1 (Type I_n — Dihedral groups). The *dihedral group* of isometries of a regular n -gon is a Coxeter group denoted I_n . See Figure 1 for illustrations.

Example 2.2 (Type A — Symmetric groups). The symmetric group $\mathfrak{S}_{n+1}$, acting on the hyperplane $\mathbb{1}^\perp := \{x \in \mathbb{R}^{n+1} \mid \langle \mathbb{1} | x \rangle = 0\}$ by permutation of the coordinates, is the reflection group of *type* A_n . It is the group of isometries of the standard n -dimensional regular simplex $\text{conv}\{e_1, \dots, e_{n+1}\}$. Its reflections are the transpositions of $\mathfrak{S}_{n+1}$, and we can choose the adjacent transpositions $\tau_p := (p, p+1)$ to be the simple reflections. A root system is given by $\Phi := \{e_p - e_q \mid p \neq q \in [n+1]\}$. Its set of simple roots is $\Delta := \{e_{p+1} - e_p \mid p \in [n]\}$, and its set of fundamental weights is $\nabla := \left\{ \sum_{q>p} e_q \mid p \in [n] \right\}$. Note that we have chosen these weights to match usual conventions, even if they do not live in the hyperplane $\mathbb{1}^\perp$. For example,




FIGURE 1. The dihedral groups I_4 and I_5 and their permutahedra. The fundamental chamber is the topmost one (colored), the simple roots are those just above the horizontal axis (in bold), and the positive roots are all roots above the horizontal axis.

$\sum \nabla = \sum_{q \in [n+1]} (q-1)e_q$ and the balanced A_n -permutahedron is the classical permutahedron obtained as the convex hull of all permutations of $\{0, \dots, n\}$, regarded as vectors in $\mathbb{R}^{n+1}$. Moreover, it enables us to match the presentation of [PS12] in the classical situation 6. The interested reader might prefer to project the weights to the hyperplane $\mathbb{1}^\perp$, and adapt the examples in type A accordingly. Figure 2 presents the A_3 -arrangement and a A_3 -permutahedron, where we marked the bottommost vertex with a dot to distinguished which chamber will be considered to be the fundamental chamber in further pictures.

Example 2.3 (Type B_n — Hyperoctahedral groups). Consider the symmetry group of $\mathfrak{S}_n$ acting on $\mathbb{R}^n$ by permutation of the coordinates together with the group $(\mathbb{Z}_2)^n$ acting on $\mathbb{R}^n$ by sign change. The semidirect product $B_n := \mathfrak{S}_n \rtimes (\mathbb{Z}_2)^n$ given by this action is a reflection group. It is the isometry group of the n -dimensional regular cross-polytope $\text{conv}\{\pm e_1, \dots, \pm e_n\}$ and of its polar n -dimensional regular cube $[-1, 1]^n$. Its reflections are the transpositions of $\mathfrak{S}_n$ and the changes of one sign. A root system is given by $\Phi := \{\pm e_p \pm e_q \mid p < q \in [n]\} \cup \{\pm e_p \mid p \in [n]\}$, and we can choose $\Delta := \{e_1\} \cup \{e_{p+1} - e_p \mid p \in [n-1]\}$ for the set of simple roots. Figure 3 shows the B_3 -arrangement and a B_3 -permutahedron, where we marked the bottommost vertex with a dot to distinguished the fundamental chamber.

Example 2.4 (Type H_3 — Icosahedral group). The isometry group of the regular icosahedron (and of its polar regular dodecahedron) is a reflection group. It is isomorphic to the direct product of $\mathbb{Z}_2$ by the alternating group $\mathfrak{A}_5$ of even permutations of $\{1, \dots, 5\}$. Figure 4 presents the H_3 -arrangement and a H_3 -permutahedron, where the fundamental chamber is marked by a dot.

2.2. Subword complexes. Let (W, S) be a Coxeter system, let $Q := q_1 \cdots q_m \in S^*$ and let $\rho \in W$. In [KM04, KM05], A. Knutson and E. Miller define the *subword complex* $\mathcal{S}(Q, \rho)$ as the pure simplicial complex of subwords of Q whose complements contain a reduced expression of ρ . The vertices of this simplicial complex are labeled by (positions of) the letters in the word Q . Note that two positions are different even if the letters of Q at these positions coincide. We denote by $[m] := \{1, \dots, m\}$




FIGURE 2. The A_3 -arrangement, intersected with the ball (left) and with the corresponding two polar regular tetrahedra (middle). The A_3 -permutahedron is a *truncated octahedron* (right).




FIGURE 3. The B_3 -arrangement, intersected with the ball (left) and with the regular cube and the regular octahedron (middle). The B_3 -permutahedron is a *great rhombicuboctahedron* (right).




FIGURE 4. The H_3 -arrangement, intersected with the ball (left) and with the regular icosahedron and the regular dodecahedron (middle). The H_3 -permutahedron is a *great rhombicosidodecahedron* (right).

the set of positions in Q . The facets of the subword complex $\mathcal{S}(Q, \rho)$ are the complements of the reduced expressions of ρ in the word Q .

It is proved in [KM04] that the subword complex $\mathcal{S}(Q, \rho)$ is a topological sphere if $\rho = \delta(Q)$, and a ball otherwise. In this paper we only consider spherical subword complexes. By appending to the word Q any reduced expression of $\delta(Q)^{-1}w_\circ$, we obtain a word Q' whose Demazure product is $\delta(Q') = w_\circ$ and such that the spherical subword complexes $\mathcal{S}(Q, \delta(Q))$ and $\mathcal{S}(Q', w_\circ)$ are isomorphic (see [CLS11, Theorem 3.7] for details). Thus, we assume that $\rho = \delta(Q) = w_\circ$ to simplify the presentation without loss of generality, and we write $\mathcal{S}(Q)$ instead of $\mathcal{S}(Q, w_\circ)$ to simplify notations.

For any facet I of $\mathcal{S}(Q)$ and any element $i \in I$, there is a unique facet J of $\mathcal{S}(Q)$ and a unique element $j \in J$ such that $I \setminus i = J \setminus j$. We say that I and J are *adjacent* facets, and that J is obtained from I by *flipping* i .

Example 2.5 (A toy example). To illustrate definitions and results throughout the paper, we will follow all details of the example of the word $Q^{\text{ex}} := \tau_2\tau_3\tau_1\tau_3\tau_2\tau_1\tau_2\tau_3\tau_1$ whose letters are adjacent transpositions of $\mathfrak{S}_4$. The facets of the subword complex $\mathcal{S}(Q^{\text{ex}})$ are $\{2, 3, 5\}$, $\{2, 3, 9\}$, $\{2, 5, 6\}$, $\{2, 6, 7\}$, $\{2, 7, 9\}$, $\{3, 4, 5\}$, $\{3, 4, 9\}$, $\{4, 5, 6\}$, $\{4, 6, 7\}$, and $\{4, 7, 9\}$.

The classical situation 1 (Sorting networks). Let $W := \mathfrak{S}_{n+1}$ act on $\mathbb{1}^\perp$ and $S := \{\tau_p \mid p \in [n]\}$, where τ_p denotes the adjacent transposition $(p, p+1)$. We can represent the word $Q := q_1q_2 \cdots q_m \in S^*$ by a *sorting network* $\mathcal{N}_Q$ as illustrated in Figure 5 (left). The network $\mathcal{N}_Q$ is formed by $n+1$ horizontal lines (its *levels*, labeled from bottom to top) together with m vertical segments (its *commutators*, labeled from left to right) corresponding to the letters of Q . If $q_k = \tau_p$, the k^{th} commutator of $\mathcal{N}_Q$ lies between the p^{th} and $(p+1)^{\text{th}}$ levels of $\mathcal{N}_Q$.




FIGURE 5. The sorting network $\mathcal{N}_{Q^{\text{ex}}}$ corresponding to the word Q^{ex} of Example 2.5 (left) and the pseudoline arrangements supported by the network $\mathcal{N}_{Q^{\text{ex}}}$ corresponding to the adjacent facets $\{2, 3, 5\}$ (middle) and $\{2, 3, 9\}$ (right) of $\mathcal{S}(Q^{\text{ex}})$.

A *pseudoline* supported by $\mathcal{N}_Q$ is an abscissa monotone path on the network $\mathcal{N}_Q$. A commutator of $\mathcal{N}_Q$ is a *crossing* between two pseudolines if it is traversed by both pseudolines, and a *contact* if its endpoints are contained one in each pseudoline. A *pseudoline arrangement* Λ (with contacts) is a set of n pseudolines on $\mathcal{N}_Q$, any two of which have precisely one crossing, possibly some contacts, and no other intersection. As a consequence of the definition, the pseudoline of Λ which starts at level p ends at level $n - p + 1$, and is called the p^{th} pseudoline of Λ . As illustrated in Figure 5 (middle and right), a facet I of $\mathcal{S}(Q)$ is represented by a pseudoline arrangement Λ_I supported by $\mathcal{N}_Q$. Its contacts (resp. crossings) are the commutators of $\mathcal{N}_Q$ corresponding to the letters of I (resp. of the complement of I).

Let I and J be two adjacent facets of $\mathcal{S}(Q)$, with $I \setminus i = J \setminus j$. Then j is the position of the crossing between the two pseudolines of Λ_I which are in contact

at position i , and the pseudoline arrangement Λ_J is obtained from the pseudoline arrangement Λ_I by exchanging the contact at i with the crossing at j . The flip between the pseudoline arrangements associated to two adjacent facets of $\mathcal{S}(Q^{\text{ex}})$ is illustrated in Figure 5 (middle and right).

A *brick* of $\mathcal{N}_Q$ is a connected component of its complement, bounded on the right by a commutator of $\mathcal{N}_Q$. For $k \in [m]$, the k^{th} brick is that immediately to the left of the k^{th} commutator of $\mathcal{N}_Q$.

The classical situation 2 (Combinatorial model for families of geometric graphs). As pointed out in [Stu11, PP12], type A subword complexes can be used to provide a combinatorial model for relevant families of geometric graphs. It relies on the interpretation of these geometric graphs in the line space of the plane. We do not present this interpretation and refer to [PP12] for details. As motivation, we just recall the following four families of geometric graphs which can be interpreted in terms of sorting networks (see Figure 6 for illustrations).

- (i) Let c be a reduced expression of a Coxeter element c and $w_o(c)$ denotes the c -sorting word of w_o (see Section 6 for definitions). Then the subword complex $\mathcal{S}(cw_o(c))$ is isomorphic to the simplicial complex of crossing-free sets of internal diagonals of the $(n+3)$ -gon, whose facets are triangulations of the $(n+3)$ -gon. It is the boundary complex of the polar of the *associahedron*.
- (ii) The subword complex $\mathcal{S}(c^k w_o(c))$ is isomorphic to the simplicial complex of $(k+1)$ -crossing free sets of k -relevant diagonals of the $(n+2k+1)$ -gon, whose facets are k -triangulations of the $(n+2k+1)$ -gon [PS09, PP12, Stu11].
- (iii) The simplicial complex of pointed crossing-free sets of internal edges of a set P of $n+3$ points in the plane, whose facets are pseudotriangulations of P [PV96, RSS08], is isomorphic to a type A_n subword complex $\mathcal{S}(Q)$, where Q is obtained from the dual pseudoline arrangement of P by removing its first and last levels [PP12]. When P is an euclidean planar point set, this




FIGURE 6. Sorting networks interpretations of certain geometric graphs: a triangulation of the convex octagon, a 2-triangulation of the convex octagon, a pseudotriangulation of a point set, and a pseudotriangulation of a set of disjoint convex bodies.

simplicial complex is known to be the boundary complex of the polar of the *pseudotriangulation polytope* of [RSS03].

- (iv) For odd n , the simplicial complex of crossing-free sets of internal free bitangents of a set X of $(n+3)/2$ disjoint convex bodies of the plane, whose facets are pseudotriangulations of X [PV96], is isomorphic to a type A_n subword complex $\mathcal{S}(Q)$, where Q is obtained from the dual double pseudoline arrangement of X by removing its first and last levels [PP12].

The classical situation 3 (Symmetric sorting networks). Let $W := B_n$ and let $S := \{\chi\} \cup \{\tau_p \mid p \in [n-1]\}$, where χ denotes the operator which changes the sign of the first coordinate and τ_p denotes the adjacent transposition $(p, p+1)$. If the word $Q \in S^*$ has x occurrences of χ , then it can be interpreted as a horizontally symmetric network $\mathcal{N}_Q$ with $2n+2$ levels and $2m-x$ commutators: replace each appearance of χ by a commutator between the first levels above and below the axis, and each appearance of τ_p by a symmetric pair of commutators between the p^{th} and $(p+1)^{\text{th}}$ levels both above and below the axis. The reduced expressions of $w_\circ$ in Q correspond to the horizontally symmetric pseudoline arrangements supported by $\mathcal{N}_Q$. Since the horizontal symmetry in the dual space corresponds to the central symmetry in the primal space, the centrally symmetric versions of the geometric objects discussed in the classical situation 2 can be interpreted combinatorially by well-chosen type B subword complexes.

3. ROOT CONFIGURATIONS

Definition 3.1. To a facet I of $\mathcal{S}(Q)$ and a position $k \in [m]$, we associate the root

$$r(I, k) := \sigma_I^{[1, k-1]}(\alpha_{q_k}),$$

where $\sigma_I^{[x, y]}$ denotes the product of the reflections q_z for $z \in [x, y] \setminus I$ in this order. The *root configuration* of the facet I is the multiset of all roots associated to the elements of I ,

$$R(I) := \{r(I, i) \mid i \in I\}.$$

Example 3.2. In the subword complex $\mathcal{S}(Q^{\text{ex}})$ with $Q^{\text{ex}} = \tau_2\tau_3\tau_1\tau_3\tau_2\tau_1\tau_2\tau_3\tau_1$ of Example 2.5, we have for example $r(\{2, 3, 9\}, 2) = \tau_2(e_4 - e_3) = e_4 - e_2$ and $r(\{2, 3, 9\}, 7) = \tau_2\tau_3\tau_2\tau_1(e_3 - e_2) = e_3 - e_1$. Moreover,

$$R(\{2, 3, 9\}) = \{e_4 - e_2, e_3 - e_1, e_3 - e_4\}.$$

The classical situation 4 (Incident pseudolines). Keep the notations of the classical situation 1. For any $k \in [m]$, we have $r(I, k) = e_t - e_b$, where t and b are such that the t^{th} and b^{th} pseudolines of Λ_I arrive respectively on top and bottom of the k^{th} commutator of $\mathcal{N}_Q$. For example, in the subword complex of Example 2.5, the root $r(\{2, 3, 9\}, 7) = e_3 - e_1$ can be read in the 7^{th} commutator of Figure 5 (right).

In [PS12], the root configuration $R(I)$ is studied as the incidence configuration of the contact graph $\Lambda_I^\#$ of the pseudoline arrangement Λ_I . This oriented graph has one node for each pseudoline of Λ_I and one arc for each contact of Λ_I , oriented from the pseudoline passing above to the pseudoline passing below. We avoid this notion here since it is not needed for the construction of this paper, and only provides a graphical description in type A which does not generalize to general finite types.

Throughout this paper, we show that the combinatorial and geometric properties of the root configuration $R(I)$ encode many relevant properties of the facet I . We

first note that, for a given facet I of $\mathcal{S}(\mathbf{Q})$, the map $\mathbf{r}(I, \cdot) : [m] \rightarrow \Phi$ can be used to understand the flips in I . We refer to Figure 5 for an illustration of the properties of the next lemma in type A. It can also be found in [CLS11, Section 3.1].

Lemma 3.3. *Let I be any facet of the subword complex $\mathcal{S}(\mathbf{Q})$.*

- (1) *The map $\mathbf{r}(I, \cdot) : k \mapsto \mathbf{r}(I, k)$ is a bijection between the complement of I and Φ^+ .*
- (2) *If I and J are two adjacent facets of $\mathcal{S}(\mathbf{Q})$ with $I \setminus i = J \setminus j$, the position j is the unique position in the complement of I for which $\mathbf{r}(I, j) \in \{\pm \mathbf{r}(I, i)\}$. Moreover, $\mathbf{r}(I, j) = \mathbf{r}(I, i) \in \Phi^+$ if $i < j$, while $\mathbf{r}(I, j) = -\mathbf{r}(I, i) \in \Phi^-$ if $j < i$.*
- (3) *In the situation of (2), the map $\mathbf{r}(J, \cdot)$ is obtained from the map $\mathbf{r}(I, \cdot)$ by:*

$$\mathbf{r}(J, k) = \begin{cases} s_{\mathbf{r}(I, i)}(\mathbf{r}(I, k)) & \text{if } \min(i, j) < k \leq \max(i, j), \\ \mathbf{r}(I, k) & \text{otherwise.} \end{cases}$$

Proof. Choose a positive root $\beta \in \Phi^+$. Since $\sigma_I^{[1, m]}$ is a reduced expression for $w_\circ$, we know that $(\sigma_I^{[1, m]})^{-1}(\beta) = w_\circ(\beta) \in \Phi^-$. Consequently, there exists $k \in [m]$ such that $(\sigma_I^{[1, k-1]})^{-1}(\beta) \in \Phi^+$ while $q_k(\sigma_I^{[1, k-1]})^{-1}(\beta) \in \Phi^-$. Therefore, we have that $\alpha_{q_k} = (\sigma_I^{[1, k-1]})^{-1}(\beta)$, and thus $\beta = \mathbf{r}(I, k)$. This proves that $\mathbf{r}(I, \cdot)$ is surjective, and even bijective since Φ^+ and the complement of I have the same cardinality.

For (3), assume that $i < j$. Since $\mathbf{r}(I, k) = \sigma_I^{[1, k-1]}(\alpha_{q_k}) = -(\sigma_I^{[k, m]})^{-1}(\alpha_{q_k})$, the roots $\mathbf{r}(I, k)$ and $\mathbf{r}(J, k)$ are equal as soon as I and J coincide either on $\{1, \dots, k-1\}$ or on $\{k, \dots, m\}$. This proves (3) when $k \leq i$ or $j < k$. In the case where $i < k \leq j$, we have

$$\mathbf{r}(J, k) = \sigma_I^{[1, i-1]} q_i \sigma_I^{[i, k-1]}(\alpha_{q_k}) = s_{\sigma_I^{[1, i-1]}(\alpha_{q_i})} \sigma_I^{[1, i-1]} \sigma_I^{[i, k-1]}(\alpha_{q_k}) = s_{\mathbf{r}(I, i)}(\mathbf{r}(I, k)),$$

where the second equality comes from the commutation rule $ws_v = s_{w(v)}w$. The proof is similar if $j < i$.

Finally, (2) is a direct consequence of (1) and (3): when we remove i from I , we have to add a position $j \notin I$ such that the root function keeps mapping the complement of $(I \setminus i) \cup j$ to Φ^+ . $\square$

Lemma 3.4. *The multiset $\mathbf{R}(I)$ characterizes I among all facets of $\mathcal{S}(\mathbf{Q})$.*

Proof. Let I and J be two distinct facets of $\mathcal{S}(\mathbf{Q})$. Assume without loss of generality that the first element i of the symmetric difference $I \triangle J$ is in $I \setminus J$. Since $\mathbf{r}(I, i)$ only depends on $I \cap [i-1]$, the roots $\mathbf{r}(I, k)$ and $\mathbf{r}(J, k)$ coincide for all $1 \leq k \leq i$. Let $\alpha := \sigma_I^{[1, i-1]}(\alpha_{q_i}) = \sigma_J^{[1, i-1]}(\alpha_{q_i}) \in \Phi^+$. According to Lemma 3.3(2), all occurrences of α in $\mathbf{R}(J)$ appear before i . Consequently, α appears at least once more in $\mathbf{R}(I)$ than in $\mathbf{R}(J)$. $\square$

Remark 3.5. Lemma 3.4 says that we can reconstruct the facet I from the root configuration $\mathbf{R}(I)$ scanning the word $\mathbf{Q}$ from left to right as follows. We start from position 0 in $\mathbf{Q}$ and we define I_0 to be the empty word and $\mathbf{R}_0$ to be the multiset $\mathbf{R}(I)$. At step k , we construct a new subword I_k from I_{k-1} and a new multiset $\mathbf{R}_k$ from $\mathbf{R}_{k-1}$ as follows:

- if $\sigma_{I_{k-1}}^{[1, k-1]}(\alpha_{q_k}) \in \mathbf{R}_{k-1}$, then $I_k := I_{k-1} \cup k$ and $\mathbf{R}_k := \mathbf{R}_{k-1} \setminus \sigma_{I_{k-1}}^{[1, k-1]}(\alpha_{q_k})$;
- otherwise, $I_k := I_{k-1}$ and $\mathbf{R}_k := \mathbf{R}_{k-1}$.

The facet I is the subword I_m we obtained at the end of this procedure. Observe that we can even reconstruct I knowing only its positive roots $\mathbf{R}(I) \cap \Phi^+$. Indeed, in

the previous sweeping procedure, the positions k for which $\sigma_{I_{k-1}}^{[1,k-1]}(\alpha_{q_k})$ is a negative root are forced to belong to the facet I according to Lemma 3.3(1). Similarly, we can reconstruct the facet I knowing only its negative roots $R(I) \cap \Phi^-$, sweeping the word Q from right to left.

The geometry of the root configuration $R(I)$ encodes many combinatorial properties of I . In fact, the facet I is relevant for the brick polytope only when its root configuration is pointed. This is in particular the case when $R(I)$ forms a basis of V . From Lemma 3.3 and the connectedness of the flip graph, we derive that this property depends only on Q , not on the particular facet I of $\mathcal{S}(Q)$.

Lemma 3.6. *Either all the root configurations $R(I)$, for I facet of $\mathcal{S}(Q)$, are simultaneously linear bases of the vector space V , or none of them is.*

In this paper, we assume that the word Q is such that the root configuration $R(I)$ forms a linear basis of V for any facet I of $\mathcal{S}(Q)$. We say that the word Q is *realizing*, and we denote by $\mathfrak{R}$ the class of all realizing words. The following proposition ensures that if the root configuration is linearly independent but does not span V , then we can always find an isomorphic subword complex for a parabolic subgroup of W for which the word is indeed realizing.

Proposition 3.7. *Let Q be a word such that the root configuration $R(I)$ is linearly independent. Then there exists a word Q' for a parabolic subgroup of W such that $\mathcal{S}(Q)$ and $\mathcal{S}(Q')$ are isomorphic and moreover, that $R(I')$ forms indeed a linear basis for any facet I' of $\mathcal{S}(Q')$.*

Proof. The proof follows exactly the same lines as that of [CLS11, Lemma 5.4]. We write here $r_Q(I, k)$ and $R_Q(I)$ to specify that we consider the root function and the root configuration on the underlying word Q . Consider a facet I of $\mathcal{S}(Q)$. Let W' be the parabolic subgroup of W generated by the reflections $\{s_{r_Q(I, i)} \mid i \in I\}$, let S' be the generators of S in W' , and let Φ' be the corresponding subsystem of roots. Let $P' := \{k \in [m] \mid r_Q(I, k) \in \Phi'\}$, and $m' := |P'|$. We define a bijection $\phi : P' \rightarrow [m']$ by $\phi(k) := |\{r_Q(I, j) \mid j \leq k\} \cap \Phi'|$. Then there is a unique word Q' on S' such that $r_Q(I, k) = r_{Q'}(\phi(I), \phi(k))$, for any $k \in P'$, which can be constructed scanning any facet I of $\mathcal{S}(Q)$ from left to right. This word Q' clearly has the desired properties. $\square$

The classical situation 5 (Contact tree). In type A , the root configuration $R(I)$ forms a basis of $\mathbb{I}^\perp$ if and only if the contact graph of Λ_I is a tree. It is independent if and only if the contact graph of Λ_I is a forest. For example, the word Q^{ex} of Example 2.5 is realizing.

Example 3.8 (Duplicated word). Let $w_\circ := w_1 \cdots w_N$ be a reduced expression of the longest element $w_\circ$ of W (thus $N = \ell(w_\circ) = |\Phi^+|$). For $k \in [N]$, we define a root $\alpha_k := w_1 \cdots w_{k-1}(\alpha_{w_k})$. Let P be a set of n positions in $w_\circ$ such that the corresponding roots $\{\alpha_p \mid p \in P\}$ form a linear basis of V . Define a new word $Q^{\text{dup}} \in S^*$ obtained by duplicating the letters of $w_\circ$ at positions in P . For $k \in [N]$, let $k^* := k + |P \cap [k-1]|$ be the new position in Q^{dup} of the k^{th} letter of $w_\circ$. Thus, we have $[N+n] = \{k^* \mid k \in [N]\} \sqcup \{p^* + 1 \mid p \in P\}$. The facets of $\mathcal{S}(Q^{\text{dup}})$ are precisely the sets $I_\varepsilon := \{p^* + \varepsilon_p \mid p \in P\}$ where $\varepsilon := (\varepsilon_1, \dots, \varepsilon_n) \in \{0, 1\}^P$. Thus, the subword complex $\mathcal{S}(Q^{\text{dup}})$ is the boundary complex of the n -dimensional cross polytope. Moreover, the roots of a facet I_ε of $\mathcal{S}(Q^{\text{dup}})$ are given by $r(I_\varepsilon, k^*) = \alpha_k$

for $k \in [N]$ and $r(I_\varepsilon, p^* + 1) = (-1)^{\varepsilon_p} \alpha_p$ for $p \in P$. Thus, the root configuration of I_ε is given by $R(I_\varepsilon) = \{(-1)^{\varepsilon_p} \alpha_p \mid p \in P\}$, and the word Q^{dup} is realizing. To illustrate the results of this paper, we discuss further the properties of duplicated words in Examples 4.3, 5.2, 5.7, 5.10, 5.18 and 5.21.

We now consider the geometry of the root configuration $R(I)$ of a facet I of $\mathcal{S}(Q)$ with respect to a linear functional $f : V \rightarrow \mathbb{R}$. We say that the flip of an element i in I is *f-preserving* if $f(r(I, i)) = 0$. We denote by

$$\mathcal{S}_f(Q) := \{I \text{ facet of } \mathcal{S}(Q) \mid \forall i \in I, f(r(I, i)) \geq 0\}$$

the set of facets of $\mathcal{S}(Q)$ whose root configuration is contained in the closed positive halfspace defined by f .

Proposition 3.9. *The set $\mathcal{S}_f(Q)$ forms a connected component of the graph of f -preserving flips on $\mathcal{S}(Q)$.*

Proof. Assume that I and J are two adjacent facets of $\mathcal{S}(Q)$ related by an f -preserving flip. Write $I \setminus i = J \setminus j$ with $f(r(I, i)) = 0$. By Lemma 3.3(3), we have $r(J, k) - r(I, k) \in \mathbb{R} \cdot r(I, i)$ and thus $f(r(J, k)) = f(r(I, k))$ for all $k \in [m]$. Thus, $\mathcal{S}_f(Q)$ is closed under f -preserving flips.

It remains to prove that $\mathcal{S}_f(Q)$ is connected by f -preserving flips. Let I be a facet of $\mathcal{S}_f(Q)$, and define $P := \{i \in I \mid f(r(I, i)) > 0\}$. We prove that we can reconstruct P , using an inductive argument on the cardinality of P . The connectedness of $\mathcal{S}_f(Q)$ then follows from that of the subword complex $\mathcal{S}(Q_{[m] \setminus P})$ of the word obtained from Q by erasing the letters at the positions given by P .

Assume first that $P = \{p\}$, and that $r(I, p) \in \Phi^-$. Performing f -preserving flips, we can make sure that $r(I, i) \in \Phi^-$ for each $i \in I$ with $f(r(I, i)) = 0$. Then $j \in I$ if and only if $\sigma_{I \cap [j-1]}^{[1, j-1]}(\alpha_{q_j}) \in f^{-1}(\mathbb{R}_{>0}) \cap \Phi^-$, and thus the facet I can be reconstructed scanning the word Q from left to right. If $r(I, p) \in \Phi^+$, we argue similarly scanning the word Q from right to left.

Assume now that $P = P' \cup \{p\}$, with $P' \neq \emptyset$. Consider an auxiliary linear function f' vanishing on $R(I) \cap f^{-1}(0)$ and on $r(I, p)$ (this function exists because $R(I)$ forms a basis of V). Then I is a facet of $\mathcal{S}_{f'}(Q)$, so that we can reconstruct the positions P' by induction hypothesis. Finally, since $I \setminus P'$ is also a facet of $\mathcal{S}(Q_{[m] \setminus P'})$, we can recover the last position p with a similar argument as before. $\square$

Corollary 3.10. *The set $\mathcal{S}_f(Q)$ is a face of the subword complex $\mathcal{S}(Q)$.*

Proof. Define again the set of positions $P := \{k \in [m] \mid f(r(I, k)) > 0\}$ for some facet I of $\mathcal{S}_f(Q)$. Since the set $\mathcal{S}_f(Q)$ is connected by f -preserving flips, which preserve the value $f(r(I, k))$ for all $k \in [m]$, the set P is independent of the choice of $I \in \mathcal{S}_f(Q)$. Furthermore, any position $i \in I \setminus P$ is an f -preserving flip. Consequently, $\mathcal{S}_f(Q)$ is precisely the set of facets of $\mathcal{S}(Q)$ which contains P . $\square$

4. GENERALIZED BRICK POLYTOPES

We begin this section with generalizing the definition of brick polytopes of [PS12] to any spherical subword complex $\mathcal{S}(Q)$, for $Q := q_1 q_2 \cdots q_m \in S^*$ with $\delta(Q) = w_\circ$.

Definition 4.1. *To a facet I of $\mathcal{S}(Q)$ and a position $k \in [m]$, we associate the weight*

$$w(I, k) := \sigma_I^{[1, k-1]}(\omega_{q_k}),$$

where $\sigma_I^{[x,y]}$ denotes the product of the reflections q_z for $z \in [x, y] \setminus I$ in this order. The **brick vector** of the facet I is the vector

$$\mathbf{B}(I) := \sum_{k \in [m]} \mathbf{w}(I, k).$$

The **brick polytope** $\mathcal{B}(\mathbf{Q})$ of the word $\mathbf{Q}$ is the convex hull of all the brick vectors, $\mathcal{B}(\mathbf{Q}) := \text{conv} \{ \mathbf{B}(I) \mid I \text{ facet of } \mathcal{S}(\mathbf{Q}) \}$.

Example 4.2. In the subword complex $\mathcal{S}(\mathbf{Q}^{\text{ex}})$ of Example 2.5, we have for example $\mathbf{w}(\{2, 3, 9\}, 2) = \tau_2(e_4) = e_4$ and $\mathbf{w}(\{2, 3, 9\}, 7) = \tau_2\tau_3\tau_2\tau_1(e_3 + e_4) = e_2 + e_3$. The brick vector of $\{2, 3, 9\}$ is $\mathbf{B}(\{2, 3, 9\}) = e_1 + 6e_2 + 5e_3 + 6e_4 = (1, 6, 5, 6)$. The brick polytope $\mathcal{B}(\mathbf{Q}^{\text{ex}})$ is a pentagonal prism, represented in Figure 7.




FIGURE 7. The brick polytope $\mathcal{B}(\mathbf{Q}^{\text{ex}})$ of the word $\mathbf{Q}^{\text{ex}}$ of Example 2.5.

The classical situation 6 (Counting bricks). In type A , we can read this definition on the sorting network interpretation of the classical situation 1. For a facet I of $\mathcal{S}(\mathbf{Q})$ and a position $k \in [m]$, the weight $\mathbf{w}(I, k)$ is the characteristic vector of the pseudolines of Λ_I which pass above the k^{th} brick of $\mathcal{N}_{\mathbf{Q}}$. For any $p \in [n+1]$, the p^{th} coordinate of the brick vector $\mathbf{B}(I)$ is the number of bricks of $\mathcal{N}_{\mathbf{Q}}$ below the p^{th} pseudoline of Λ_I . See Figure 5 for an illustration. This was the original definition used in type A in [PS12], which explains the name **brick polytope**.

Example 4.3 (Duplicated word). Consider the situation of Example 3.8: the word $\mathbf{Q}^{\text{dup}}$ is obtained from the reduced expression $w_{\circ} := w_1 \cdots w_N$ of $w_{\circ}$ by duplicating each letter located at a position in the chosen set P . For $k \in [N]$, we define a weight $\omega_k := w_1 \cdots w_{k-1}(\omega_{w_k})$. Then the weights of a facet I_{ε} of $\mathcal{S}(\mathbf{Q}^{\text{dup}})$ are given by $\mathbf{w}(I_{\varepsilon}, k^*) = \omega_k$ for $k \in [N]$ and $\mathbf{w}(I_{\varepsilon}, p^* + 1) = \omega_p + \varepsilon_p \alpha_p$ for $p \in P$. Consequently, the brick vector of the facet I_{ε} is given by $\mathbf{B}(I_{\varepsilon}) = \Theta + \sum_{p \in P} \varepsilon_p \alpha_p$, where $\Theta := \sum_{k \in [N]} \omega_k + \sum_{p \in P} \omega_p$. It follows that the brick polytope $\mathcal{B}(\mathbf{Q}^{\text{dup}})$ is an n -dimensional parallelepiped, whose edges are directed by the basis $\{\alpha_p \mid p \in P\}$.

Although we define brick polytopes for any word $\mathbf{Q}$, we only present in this paper their geometric and combinatorial properties when $\mathbf{Q}$ is realizing. We will see that

this condition is sufficient for the brick polytope to realize its subword complex. Reciprocally, the brick polytope can realize its subword complex only when the root configurations of the facets are linearly independent, and we can then assume that the word Q is realizing by Proposition 3.7. A remark on the situation of non-realizing words can be found in the conclusion.

The root function $r(\cdot, \cdot)$ and the weight function $w(\cdot, \cdot)$ have similar definitions and are indeed closely related. The following lemma underlines some of their similarities.

Lemma 4.4. *Let I be a facet of the subword complex $\mathcal{S}(Q)$.*

- (1) *We have $w(I, k+1) = \begin{cases} w(I, k) & \text{if } k \in I, \\ w(I, k) - r(I, k) & \text{if } k \notin I. \end{cases}$*
- (2) *If I and J are two adjacent facets of $\mathcal{S}(Q)$, with $I \setminus i = J \setminus j$, then $w(J, \cdot)$ is obtained from $w(I, \cdot)$ by:*

$$w(J, k) = \begin{cases} s_{r(I, i)}(w(I, k)) & \text{if } \min(i, j) < k \leq \max(i, j), \\ w(I, k) & \text{otherwise.} \end{cases}$$

- (3) *For $j \notin I$, we have $\langle r(I, j) | w(I, k) \rangle$ is non-negative if $j \geq k$, and non-positive if $j < k$.*

Proof. (1) is an immediate consequence of the definitions of $r(I, k)$ and $w(I, k)$. The proof of (2) is similar to the proof of Lemma 3.3(2). We now prove (3). If $j \geq k$, then $\sigma_I^{[k, j-1]}(\alpha_{q_j}) \in \Phi^+$ since $j \notin I$. Thus, $\langle r(I, j) | w(I, k) \rangle = \langle \sigma_I^{[k, j-1]}(\alpha_{q_j}) | \omega_{q_k} \rangle \geq 0$. Similarly, if $j < k$, then $(\sigma_I^{[j, k-1]})^{-1}(\alpha_{q_j}) \in \Phi^-$ since $j \notin I$. Consequently, $\langle r(I, j) | w(I, k) \rangle = \langle (\sigma_I^{[j, k-1]})^{-1}(\alpha_{q_j}) | \omega_{q_k} \rangle \leq 0$. $\square$

Lemma 4.5. *If I and J are two adjacent facets of $\mathcal{S}(Q)$, with $I \setminus i = J \setminus j$, then the difference of the brick vectors $B(I) - B(J)$ is a positive multiple of $r(I, i)$.*

Proof. Since I and J play symmetric roles, we can assume that $i < j$. Lemma 4.4 implies that

$$B(I) - B(J) = \sum_{k \in [m]} w(I, k) - w(J, k) = \left(\sum_{i < k \leq j} 2 \langle r(I, j) | w(I, k) \rangle \right) \cdot r(I, i),$$

where the last sum is positive since all its summands are non-negative and the last one is positive. $\square$

Let $f : V \rightarrow \mathbb{R}$ be a linear functional. We denote by $\mathcal{B}_f(Q)$ the face of $\mathcal{B}(Q)$ which maximizes f . Remember that $\mathcal{S}_f(Q)$ denotes the set of facets of $\mathcal{S}(Q)$ whose root configuration is included in the closed positive halfspace defined by f .

Lemma 4.6. *The faces of the brick polytope $\mathcal{B}(Q)$ are faces of the subword complex $\mathcal{S}(Q)$. For any facet I of $\mathcal{S}(Q)$, we have $B(I) \in \mathcal{B}_f(Q) \iff I \in \mathcal{S}_f(Q)$.*

Proof. Let I be a facet of $\mathcal{S}(Q)$ such that $B(I) \in \mathcal{B}_f(Q)$. Let $i \in I$, let J be the facet obtained from I by flipping i , and let $\lambda > 0$ be such that $r(I, i) = \lambda(B(I) - B(J))$ (by Lemma 4.5). Then $f(r(I, i)) = \lambda(f(B(I)) - f(B(J))) \geq 0$ since $B(I)$ maximizes f . Thus, $I \in \mathcal{S}_f(Q)$. We obtained that the set of facets I of $\mathcal{S}(Q)$ such that $B(I) \in \mathcal{B}_f(Q)$ is contained in $\mathcal{S}_f(Q)$. Since the former is closed while the latter is connected under f -preserving flips (by Proposition 3.9), they coincide. $\square$

Proposition 4.7. *For any facet I of $\mathcal{S}(Q)$, the cone of the brick polytope $\mathcal{B}(Q)$ at the brick vector $B(I)$ coincides with the cone generated by the negative of the root configuration $R(I)$ of I . We set*

$$C(I) := \text{cone} \{B(J) - B(I) \mid J \text{ facet of } \mathcal{S}(Q)\} = \text{cone} \{-r(I, i) \mid i \in I\}.$$

Proof. Let $C_B(I)$ be the cone of $\mathcal{B}(Q)$ at $B(I)$ and let $C_R(I)$ be the cone generated by $-R(I)$. The inclusion $C_R(I) \subset C_B(I)$ is an immediate consequence of Lemma 4.5. For the other direction, we need to prove that any face of $C_R(I)$ is also a face of $C_B(I)$. That is to say that the brick vector $B(I)$ maximizes any linear functional f among the brick vectors of all facets of $\mathcal{S}(Q)$ as soon as it maximizes f among the brick vectors of the facets adjacent to I . This is ensured by Lemma 4.6. $\square$

Theorem 4.8. *If Q is realizing, the polar of the brick polytope $\mathcal{B}(Q)$ realizes the subword complex $\mathcal{S}(Q)$.*

Proof. Proposition 4.7 ensures that the brick vector of each facet of $\mathcal{S}(Q)$ is a vertex of $\mathcal{B}(Q)$ and that this polytope is full-dimensional and simple. Since the faces of the brick polytope are faces of the subword complex by Lemma 4.6, it immediately implies that the boundary complex of the brick polytope $\mathcal{B}(Q)$ is isomorphic to the subword complex $\mathcal{S}(Q)$. $\square$

Remark 4.9. If (W, S) is crystallographic (*i.e.* stabilizes a lattice Λ), then the brick polytope $\mathcal{B}(Q)$ is furthermore a lattice polytope (*i.e.* its vertices are elements of Λ). Indeed, the brick vector of any facet is a sum of weights of W .

5. FURTHER COMBINATORIAL AND GEOMETRIC PROPERTIES

In this section, we consider further relevant properties of the brick polytope $\mathcal{B}(Q)$. We first define a surjective map κ from the Coxeter group W to the facets of the subword complex $\mathcal{S}(Q)$. Using this map, we relate the normal fan of $\mathcal{B}(Q)$ to the Coxeter fan, and the graph of $\mathcal{B}(Q)$ to the weak order. We then study some properties of the fibers of κ with respect to the weak order. We also describe a canonical spanning tree of the graph of $\mathcal{B}(Q)$ which can be used to efficiently compute the brick vectors of all facets of $\mathcal{S}(Q)$. Finally, we decompose $\mathcal{B}(Q)$ into a Minkowski sum of W -matroid polytopes. Again, although most of the results presented in this section seem to have natural extensions to non-realizing words, their current presentation considers only realizing words $Q \in \mathfrak{R}$.

5.1. A surjective map. A set $U \subset \Phi$ is *separable* if there is a hyperplane which separates U from its complement $\Phi \setminus U$. For finite Coxeter systems, separable sets are precisely the sets $w(\Phi^+)$, for $w \in W$. According to Proposition 3.9, there is a unique facet I of $\mathcal{S}(Q)$ whose root configuration $R(I)$ is contained in a given separable set. This defines a map κ from W to the facets of $\mathcal{S}(Q)$ which associates to $w \in W$ the unique facet $\kappa(w)$ of $\mathcal{S}(Q)$ such that $R(\kappa(w)) \subset w(\Phi^+)$. Furthermore, since we assumed that any root configuration forms a basis (and thus is pointed), this map κ is surjective. However, note that this map is not injective.

Example 5.1. The map κ for the word Q^{ex} of Example 2.5 is presented in Figure 8. For each permutation $w \in \mathfrak{S}_4$, we have written the facet $\kappa(w)$ of $\mathcal{S}(Q^{\text{ex}})$, and we have represented the pseudoline arrangement $\Lambda_{\kappa(w)}$ supported by $\mathcal{N}_{Q^{\text{ex}}}$. The underlying diagram is the Hasse diagram of the weak order on $\mathfrak{S}_4$: two permutations are related by an edge if they differ by the transposition of two adjacent letters. To




FIGURE 8. The map κ from $\mathfrak{S}_4$ to the facets of the subword complex $\mathcal{S}(Q^{\text{ex}})$ of Example 2.5.

visualize the fibers of κ , we draw a light edge when the images of the permutations under κ are distinct, and a double strong edge if the images are the same.

Example 5.2 (Duplicated word). Consider the word Q^{dup} defined in Example 3.8. Two elements w, w' of W have the same image under the map κ if and only if the symmetric difference of $w(\Phi^+)$ and $w'(\Phi^+)$ is disjoint from the set $\{\pm\alpha_p \mid p \in P\}$, or equivalently the symmetric difference of the inversion sets of w and w' is disjoint from the set $\{\alpha_p \mid p \in P\}$.

Lemma 5.3. *Consider an element $w \in W$ and a simple root $\alpha \in \Delta$. Then $\kappa(ws_\alpha)$ is obtained from $\kappa(w)$ as follows.*

- (i) *If $w(\alpha) \notin R(\kappa(w))$, then $\kappa(ws_\alpha)$ equals $\kappa(w)$.*
- (ii) *Otherwise, $\kappa(ws_\alpha)$ is obtained from $\kappa(w)$ by flipping the unique $i \in \kappa(w)$ such that $w(\alpha) = r(\kappa(w), i)$.*

Proof. If $w(\alpha) \notin R(\kappa(w))$, then $R(\kappa(w)) \subset ws_\alpha(\Phi^+) = w(\Phi^+) \Delta \{\pm w(\alpha)\}$ (where Δ denotes the symmetric difference). Thus, $\kappa(ws_\alpha) = \kappa(w)$ by uniqueness in the definition of the map κ .

Assume now that $w(\alpha) = r(\kappa(w), i)$ for some $i \in \kappa(w)$, and let J denote the facet of $\mathcal{S}(Q)$ obtained from $\kappa(w)$ by flipping i , and j be the position in J but not in $\kappa(w)$. According to Lemma 3.3(2),

$$r(J, k) = \begin{cases} s_{w(\alpha)}(r(\kappa(w), k)) = ws_\alpha w^{-1}(r(\kappa(w), k)) & \text{if } \min(i, j) < k \leq \max(i, j), \\ r(\kappa(w), k) & \text{otherwise.} \end{cases}$$

If $\min(i, j) < k \leq \max(i, j)$, we have $r(J, k) \in ws_\alpha(\Phi^+)$ since $w^{-1}(r(\kappa(w), k)) \in \Phi^+$. Otherwise, we have $w^{-1}(r(\kappa(w), k)) \in \Phi^+ \setminus \alpha$ so that $s_\alpha w^{-1}(r(\kappa(w), k)) \in \Phi^+$, and thus $r(J, k) = r(\kappa(w), k) \in ws_\alpha(\Phi^+)$. We obtain that $R(J) \subset ws_\alpha(\Phi^+)$, and consequently $J = \kappa(ws_\alpha)$ by uniqueness in the definition of the map κ . $\square$

5.2. A normal fan description. Remember that the *normal cone* of a face F of a polytope $\Pi \subset V$ is the cone of all vectors $v \in V$ such that the linear function $x \mapsto \langle x|v \rangle$ on Π is maximized by all points in F . The *normal fan* of Π is the complete polyhedral fan formed by the normal cones of all faces of Π (see [Zie95, Lecture 7]). For example, the normal fan of the W -permutahedron is the Coxeter fan of W . The following proposition relates the normal fan of the brick polytope with the Coxeter fan.

Proposition 5.4. *Let I be a facet of $\mathcal{S}(Q)$. The normal cone $C^\diamond(I)$ of $\mathcal{B}(I)$ in $\mathcal{B}(Q)$ is the union of the chambers $w(\mathcal{C})$ of the Coxeter fan of W given by the elements $w \in W$ with $\kappa(w) = I$.*

Proof. For a given cone $X \subset V$, we denote by $X^\diamond := \{v \in V \mid \forall x \in X, \langle x|v \rangle \geq 0\}$ its polar cone. We have

$$C^\diamond(I) = C(I)^\diamond = \text{cone}(-R(I))^\diamond = \left(\bigcap_{w \in \kappa^{-1}(I)} w(\mathcal{C}^\diamond) \right)^\diamond = \bigcup_{w \in \kappa^{-1}(I)} w(\mathcal{C}^\diamond)^\diamond = \bigcup_{w \in \kappa^{-1}(I)} w(\mathcal{C}),$$

because $\mathcal{C}^\diamond$ is the cone generated by Φ^+ . $\square$

Corollary 5.5. *The Coxeter fan refines the normal fan of the brick polytope.*




FIGURE 9. The Coxeter fan (left) refines the normal fan of the brick polytope $\mathcal{B}(\mathcal{Q}^{\text{ex}})$ (middle) according to the fibers of κ (right).

Example 5.6. Figure 9 illustrates Corollary 5.5 for the subword complex $\mathcal{S}(\mathcal{Q}^{\text{ex}})$ of Example 2.5. The normal fan of the brick polytope $\mathcal{B}(\mathcal{Q}^{\text{ex}})$ (middle) is obtained by coarsening the chambers of the Coxeter fan (left) according to the fibers of κ (represented on the permutahedron on the right).

Example 5.7 (Duplicated word). Consider the word $\mathcal{Q}^{\text{dup}}$ defined in Example 3.8. The normal fan of its brick polytope $\mathcal{B}(\mathcal{Q}^{\text{dup}})$ is the fan formed by the hyperplanes orthogonal to the roots $\{\alpha_p \mid p \in P\}$. The vectors of the dual basis of $\{\alpha_p \mid p \in P\}$ are normal vectors for the facets of the brick polytope $\mathcal{B}(\mathcal{Q}^{\text{dup}})$. Observe that the normal fan of the brick polytope $\mathcal{B}(\mathcal{Q})$ is not in general defined by a subarrangement of the Coxeter arrangement (see *e.g.* Figure 9).

5.3. Increasing flips and quotients of the weak order. If I and J are adjacent facets of $\mathcal{S}(\mathcal{Q})$ with $I \setminus i = J \setminus j$, we say that the flip from I to J is *increasing* if $i < j$. This is equivalent to $r(I, i) \in \Phi^+$ (by Lemma 3.3(2)), and to $f(\mathcal{B}(I)) < f(\mathcal{B}(J))$, where f is a linear function positive on Φ^+ (by Lemma 4.5). The graph of increasing flips is thus clearly acyclic, and we call *increasing flip order* its transitive closure $\prec$. This order generalizes the Tamari order on triangulations.

Proposition 5.8. *A facet I is covered by a facet J in increasing flip order if and only if there exist $w_I \in \kappa^{-1}(I)$ and $w_J \in \kappa^{-1}(J)$ such that w_I is covered by w_J in weak order. We say that the increasing flip order is the quotient of the weak order by the map κ .*

Proof. Let $w \in W$ and $s \in S$ such that $w < ws$ in weak order. Since $w(\alpha_s) \in \Phi^+$, Lemma 5.3 ensures that either $\kappa(ws) = \kappa(w)$ or $\kappa(ws)$ is obtained from $\kappa(w)$ by an increasing flip.

Reciprocally, consider two facets I and J of $\mathcal{S}(\mathcal{Q})$ related by an increasing flip. We write $I \setminus i = J \setminus j$, with $i < j$. Their normal cones $\mathcal{C}^\circ(I)$ and $\mathcal{C}^\circ(J)$ are adjacent along a cone of codimension 1 which belongs to the hyperplane H orthogonal to $r(I, i) = r(J, j)$. Let $\mathcal{C}_I$ be a chamber of the Coxeter arrangement of W contained in $\mathcal{C}^\circ(I)$ and adjacent to H , and let $\mathcal{C}_J := s_{r(I, i)}(\mathcal{C}_I)$ denote the symmetric chamber with respect to H . Let w_I and w_J be such that $\mathcal{C}_I = w_I(\mathcal{C})$ and $\mathcal{C}_J = w_J(\mathcal{C})$. Then $w_I \in \kappa^{-1}(I)$ and $w_J \in \kappa^{-1}(J)$, and w_J covers w_I in weak order. $\square$




FIGURE 10. The Hasse diagram of the increasing flip order on the facets of the subword complex $\mathcal{S}(Q^{\text{ex}})$ of Example 2.5 is the quotient of the Hasse diagram of the weak order on $\mathfrak{S}_4$ by the fibers of the map κ represented in Figure 8.

It immediately follows from this proposition that $I \prec J$ in increasing flip order if there exists $w_I \in \kappa^{-1}(I)$ and $w_J \in \kappa^{-1}(J)$ such that $w_I < w_J$ in weak order. We conjecture that the converse also holds in general.

In his work on Cambrian lattices [Rea04, Rea06, Rea07b], N. Reading considers quotients of the weak order under lattice congruences, which fulfill sufficient conditions for the quotient to be moreover a lattice. In the context of brick polytopes, we obtain different partitions of W which are not lattice congruences in general (the fibers of κ are not necessarily intervals of the weak order, but they contain all intervals between two of their elements — see Section 5.4). We nonetheless conjecture that the resulting quotient is in general a lattice.

Example 5.9. Figure 10 represents the Hasse diagram of the increasing flip order on the facets of the subword complex $\mathcal{S}(Q^{\text{ex}})$ of Example 2.5. It is the quotient of the weak order by the fibers of the map κ presented in Figure 8.

Example 5.10 (Duplicated word). Consider the word Q^{dup} defined in Example 3.8. Then the quotient of the weak order by the fibers of κ is the boolean lattice.

5.4. Fibers of κ and the weak order. In this section, we discuss properties of the fibers of the map κ with respect to the right weak order. First, although these fibers are not always intervals (see Example 5.1), they are closed by taking intervals.

Lemma 5.11. *If two elements $\underline{w} \leq \overline{w}$ of W lie in the same fiber of κ , then the complete interval $\{w \in W \mid \underline{w} \leq w \leq \overline{w}\}$ lies in the same fiber of κ .*

Proof. As already observed in the proof of Proposition 5.8, $w < ws$ in weak order implies $\kappa(w) \leq \kappa(ws)$ in increasing flip order, for any $w \in W$. Consequently, if $\underline{w}$ and $\overline{w}$ lie in the same fiber, then so does any element w on a chain between $\underline{w}$ and $\overline{w}$, and thus in the interval $\{w \in W \mid \underline{w} \leq w \leq \overline{w}\}$. Another way to see this property will be given below by an alternative description of the map κ . $\square$

Next, we give an explicit description of the fibers of κ in terms of inversion sets of elements of W . Let $w \in W$ and $w = w_1 w_2 \cdots w_p$ be a reduced expression of w (with $w_1, w_2, \dots, w_p \in S$). The *inversion set* of w is the set

$$\text{inv}(w) := \Phi^+ \cap w(\Phi^-) = \{\alpha_{w_1}, w_1(\alpha_{w_2}), \dots, w_1 w_2 \cdots w_{p-1}(\alpha_{w_p})\} \subset \Phi^+.$$

This definition of inversion set is compatible with the right weak order, this is $w \leq w'$ if and only if $\text{inv}(w) \subseteq \text{inv}(w')$.

For finite Coxeter groups, the inversion sets of the elements of W are precisely the biclosed subsets of Φ^+ . A set $U \subset \Phi^+$ is *biclosed* if and only if both U and its complement $\Phi^+ \setminus U$ are closed under non-negative linear combinations in Φ^+ . In other words, U is biclosed if and only if $U \cup -(\Phi^+ \setminus U)$ is separable.

For a facet I of the subword complex $\mathcal{S}(Q)$, we define $R^+(I) := R(I) \cap \Phi^+$ and $R^-(I) := - (R(I) \cap \Phi^-)$. In other words, $R^+(I)$ and $R^-(I)$ are the subsets of Φ^+ such that $R(I) = R^+(I) \cup -R^-(I)$. With these notations, the preimage of I under κ is given by all $w \in W$ such that

$$R^-(I) \subset \text{inv}(w) \subset \Phi^+ \setminus R^+(I).$$

While the definition of the map κ provides a direct geometric relation between separable sets and root configurations, this alternative description provides a relation to inversion sets, and thus to the weak order. It moreover gives a criterion to characterize the fibers of κ which have a meet or a join. For a facet I of $\mathcal{S}(Q)$, define $\wedge(I)$ and $\vee(I)$ by

$$\wedge(I) := \bigcap_{w \in \kappa^{-1}(I)} \text{inv}(w) \quad \text{and} \quad \vee(I) := \bigcup_{w \in \kappa^{-1}(I)} \text{inv}(w).$$

Remark 5.12. The subsets $\wedge(I)$ and $\vee(I)$ can be computed from $R(I)$ by the following iterative procedure. Define two sequences $(\wedge_p(I))_{p \in \mathbb{N}}$ and $(\nabla_p(I))_{p \in \mathbb{N}}$ of collections of positive roots by

- (i) $\wedge_0(I) := R^-(I)$ and $\nabla_0(I) := R^+(I)$;
- (ii) the set $\wedge_{p+1}(I)$ is the set of positive roots α such that $\alpha = \lambda_1 \alpha_1 + \lambda_2 \alpha_2$ for some $\alpha_1, \alpha_2 \in \wedge_p(I)$ and some $\lambda_1, \lambda_2 \geq 0$, or such that $\lambda \alpha + \mu \beta \in \wedge_p(I)$ for some $\beta \in \nabla_p(I)$ and some $\lambda, \mu \geq 0$;
- (iii) the set $\nabla_{p+1}(I)$ is the set of positive roots β such that $\beta = \mu_1 \beta_1 + \mu_2 \beta_2$ for some $\beta_1, \beta_2 \in \nabla_p(I)$ and some $\mu_1, \mu_2 \geq 0$, or such that $\lambda \alpha + \mu \beta \in \nabla_p(I)$ for some $\alpha \in \wedge_p(I)$ and some $\lambda, \mu \geq 0$.

The sequences $(\wedge_p(I))_{p \in \mathbb{N}}$ and $(\nabla_p(I))_{p \in \mathbb{N}}$ are then stationary with limits $\wedge(I)$ and $\Phi^+ \setminus \vee(I)$.

Proposition 5.13. *The fiber $\kappa^{-1}(I)$ has a meet if and only if $\wedge(I)$ is biclosed and a join if and only if $\vee(I)$ is biclosed.*

Proof. We first observe that, $\kappa(w) = I$ if and only if $\wedge(I) \subset \text{inv}(w) \subset \vee(I)$. If $\wedge(I)$ is biclosed, it is the inversion set of an element $\underline{w} \in W$. Then $\underline{w}$ is in $\kappa^{-1}(I)$ and it is smaller than all the elements of $\kappa^{-1}(I)$ in weak order. It is thus the meet of $\kappa^{-1}(I)$. Reciprocally, if the fiber $\kappa^{-1}(I)$ has a meet, its inversion set is precisely $\wedge(I)$, which is thus biclosed. The proof is similar for the join. $\square$

Example 5.14. Consider the facet $\{2, 5, 6\}$ of the subword complex $\mathcal{S}(Q^{\text{ex}})$ of Example 2.5. Its root configuration is $R(\{2, 5, 6\}) = \{e_4 - e_2, e_4 - e_1, e_1 - e_3\}$, thus $R^+(\{2, 5, 6\}) = \{e_4 - e_2, e_4 - e_1\}$ and $R^-(\{2, 5, 6\}) = \{e_3 - e_1\}$. Its fiber is $\kappa^{-1}(\{2, 5, 6\}) = \{2314, 3124, 3214\}$ (see Figure 8). We have $\wedge(\{2, 5, 6\}) = \{e_3 - e_1\}$ and $\vee(\{2, 5, 6\}) = \{e_2 - e_1, e_3 - e_2, e_3 - e_1\}$. The fiber $\kappa^{-1}(\{2, 5, 6\})$ has no meet since $\wedge(\{2, 5, 6\})$ is not biclosed. However, its join is 3214 since $\vee(\{2, 5, 6\}) = \text{inv}(3214)$.

5.5. Canonical spanning trees of the increasing flip graph. We now describe two canonical spanning trees of the ridge graph of a subword complex constructed in [Pil12]. This construction yields an algorithm to exhaustively generate the facets of the subword complex, whose running time per facet and working space are both polynomial. We summarize here this construction since it is closely related to the topics developed in the previous sections and it can be used to compute the brick vectors of the facets of $\mathcal{S}(Q)$.

The construction of these trees proceed by induction on the length of the word Q . Consequently, we cannot restrict in this section to spherical subword complexes. We consider the subword complex $\mathcal{S}(Q, \rho)$ associated to a word $Q := q_1 \cdots q_m$ on S and an arbitrary element $\rho \in W$. For the induction, we define $Q_{\vdash} := q_2 \cdots q_m$ and $Q_{\dashv} := q_1 \cdots q_{m-1}$ to be the words obtained from Q erasing its first and last letters respectively. We can then decompose inductively the subword complex according on which faces contain the first or the last letter:

$$\mathcal{S}(Q, \rho) = \mathcal{S}(Q_{\vdash}, q_1 \rho)^{\rightarrow} \sqcup 1 \star \mathcal{S}(Q_{\vdash}, \rho)^{\rightarrow} = \mathcal{S}(Q_{\dashv}, \rho q_m) \sqcup \mathcal{S}(Q_{\dashv}, \rho) \star m,$$

where $\mathcal{X} \star z := z \star \mathcal{X} := \{X \cup z \mid X \in \mathcal{X}\}$ denotes the join of a set $\mathcal{X}$ of subsets of $\mathbb{N}$ with an element $z \in \mathbb{N}$, and $\mathcal{X}^{\rightarrow} := \{X^{\rightarrow} \mid X \in \mathcal{X}\} := \{\{x+1 \mid x \in X\} \mid X \in \mathcal{X}\}$ denotes the right shift of all sets in $\mathcal{X}$.

For a general subword complex $\mathcal{S}(Q, \rho)$, the increasing flip graph is again defined as the graph whose nodes are the facets of $\mathcal{S}(Q, \rho)$ and with an arc from a facet I to a facet J if and only if $I \setminus i = J \setminus j$ for some $i < j$. This graph is acyclic and has a unique source, the *positive greedy facet* $P(Q, \rho)$, and a unique sink, the *negative greedy facet* $N(Q, \rho)$. These two greedy facets are connected by the relation $N(q_1 \cdots q_m, \rho) = \{m+1-p \mid p \in P(q_m \cdots q_1, \rho^{-1})\}$. We therefore only discuss here negative greedy facets, whose properties are easier to state, and we leave to the reader the translation to positive greedy facets, which we use later in Section 6.3.2.

By definition, the negative greedy facet $N(Q, \rho)$ is the unique facet whose root configuration is a subset of negative roots, and the lexicographically largest facet of $\mathcal{S}(Q, \rho)$. Following the inductive structure of the subword complex described above, the negative greedy facet can be constructed inductively erasing the first or last letter of Q as follows:

$$N(Q, \rho) = \begin{cases} N(Q_{\vdash}, q_1 \rho)^{\rightarrow} & \text{if } \ell(q_1 \rho) < \ell(\rho) \\ 1 \cup N(Q_{\vdash}, \rho)^{\rightarrow} & \text{otherwise} \end{cases} = \begin{cases} N(Q_{\dashv}, \rho) \cup m & \text{if } \delta(Q_{\dashv}) > \rho \\ N(Q_{\dashv}, \rho q_m) & \text{otherwise} \end{cases}$$

where $X^{\rightarrow} := \{x + 1 \mid x \in X\}$ denotes the right shift of a subset $X \subset \mathbb{N}$. In other words, $N(Q, \rho)$ can be obtained sweeping the word Q either from left to right placing inversions as soon as possible, or from right to left selecting positions as long as possible. The key property of the greedy facets is the following *greedy flip property*.

Proposition 5.15 ([Pil12]). *If m is a flippable element of $N(Q, \rho)$, then $N(Q_{\rightarrow}, \rho q_m)$ is obtained from $N(Q, \rho)$ flipping m .*

From this property, we construct the *negative greedy flip tree* $\mathcal{N}(Q, \rho)$ by induction, erasing letters of Q from the right. If $Q = \varepsilon$ and $\rho = e$, then $\mathcal{N}(\varepsilon, e)$ has only one node $\emptyset$. Otherwise, we define the tree $\mathcal{N}(Q, \rho)$ to be

- $\mathcal{N}(Q_{\rightarrow}, \rho q_m)$ if m appears in none of the facets of $\mathcal{S}(Q, \rho)$;
- $\mathcal{N}(Q_{\rightarrow}, \rho) \star m$ if m appears in all the facets of $\mathcal{S}(Q, \rho)$;
- the disjoint union of $\mathcal{N}(Q_{\rightarrow}, \rho q_m)$ and $\mathcal{N}(Q_{\rightarrow}, \rho) \star m$, together with an additional arc from $N(Q_{\rightarrow}, \rho q_m)$ to $N(Q, \rho) = N(Q_{\rightarrow}, \rho) \cup m$, otherwise.

By definition, the negative greedy flip tree is a spanning tree of the increasing flip graph, oriented towards its root $N(Q, \rho)$.

The negative greedy flip tree can also be described avoiding the use of induction. Let I be a facet of $\mathcal{S}(Q, \rho)$. Define its *negative greedy index* $n(I)$ as the largest position $x \in [m]$ such that $I \cap [x] = N(q_1 \cdots q_x, \sigma_I^{[1, x]})$. This index equals m for the negative greedy facet $N(Q, \rho)$. Moreover, if I and J are two adjacent facets of $\mathcal{S}(Q, \rho)$ with $I \setminus i = J \setminus j$ and $i < j \leq n(J)$, then $n(I) = j - 1$. This index provides the following alternative induction-free description of the negative greedy flip tree.

Proposition 5.16 ([Pil12]). *The negative greedy flip tree $\mathcal{N}(Q, \rho)$ has one vertex for each facet of $\mathcal{S}(Q, \rho)$, and one arc from a facet I to a facet J if and only if $I \setminus i = J \setminus j$ for some $i \in I$ and $j \in J$ satisfying $i < j \leq n(J)$.*

The *positive greedy flip tree* $\mathcal{P}(Q, \rho)$ is defined by a similar induction, but erasing letters of Q from the left. It can also be described avoiding the use of induction. Namely, define the *positive greedy index* $p(I)$ of a facet I of $\mathcal{S}(Q, \rho)$ to be the smallest position $x \in [m]$ such that $\{i - x \mid I \setminus [x]\} = P(q_{x+1} \cdots q_m, \sigma_I^{[x+1, m]})$. This index equals 1 for the positive greedy facet $P(Q, \rho)$ and if I and J are two adjacent facets of $\mathcal{S}(Q, \rho)$ with $I \setminus i = J \setminus j$ and $p(I) \leq i < j$, then $p(J) = i + 1$. The positive greedy flip tree has one vertex for each facet of $\mathcal{S}(Q, \rho)$, and one arc from a facet I to a facet J if and only if $I \setminus i = J \setminus j$ for some $i \in I$ and $j \in J$ satisfying $p(I) \leq i < j$.

Consider now again a spherical subword complex $\mathcal{S}(Q)$ for a realizing word Q . Let f denotes a linear function positive on Φ^+ . Then the negative and positive greedy flip trees $\mathcal{N}(Q)$ and $\mathcal{P}(Q)$ are f -monotone spanning trees of the graph of the brick polytope $\mathcal{B}(Q)$, respectively rooted at the f -minimizing and f -maximizing vertices of $\mathcal{B}(Q)$. Besides this geometric interpretation, the greedy flip trees are also relevant to efficiently obtain the vertex description of the brick polytope $\mathcal{B}(Q)$. Indeed, we can compute iteratively the brick vectors of all facets of $\mathcal{S}(Q)$, propagating along the greedy flip trees.

Example 5.17. Consider the subword complex $\mathcal{S}(Q^{\text{ex}})$ of Example 2.5. Its positive greedy facet is $\{2, 3, 5\}$, while its negative greedy facet is $\{4, 7, 9\}$. The positive and negative greedy flip trees $\mathcal{P}(Q^{\text{ex}})$ and $\mathcal{N}(Q^{\text{ex}})$ of $\mathcal{S}(Q^{\text{ex}})$ are represented in Figure 11 as monotone spanning trees of the 1-skeleton of the brick polytope $\mathcal{B}(Q^{\text{ex}})$. In the positive greedy flip tree $\mathcal{P}(Q^{\text{ex}})$, each facet I is denoted as the concatenation of its

elements, where the symbol $|$ separates the elements strictly smaller than $\mathbf{p}(I)$ from those which are larger or equal to $\mathbf{p}(I)$. In the negative greedy flip tree $\mathcal{N}(\mathcal{Q}^{\text{ex}})$, each facet I is denoted as the concatenation of its elements, where the symbol $|$ separates the elements smaller or equal to $\mathbf{n}(I)$ from those which are strictly larger than $\mathbf{n}(I)$.




FIGURE 11. The positive (left) and negative (right) greedy flip trees of the subword complex $\mathcal{S}(\mathcal{Q}^{\text{ex}})$ of Example 2.5.

Example 5.18 (Duplicated word). Consider the word $\mathcal{Q}^{\text{dup}}$ defined in Example 3.8. The positive greedy facet $\mathbf{P}(\mathcal{Q}^{\text{dup}})$ is the facet $I_{(0,\dots,0)}$, while the negative greedy facet $\mathbf{N}(\mathcal{Q}^{\text{dup}})$ is the facet $I_{(1,\dots,1)}$. For any $\varepsilon := (\varepsilon_1, \dots, \varepsilon_n) \in \{0, 1\}^P$, the facet I_ε of $\mathcal{S}(\mathcal{Q}^{\text{dup}})$ has positive greedy index $\mathbf{p}(I_\varepsilon) = 1 + \max\{p^* + 1 \mid p \in P, \varepsilon_p = 1\}$ and negative greedy index $\mathbf{n}(I_\varepsilon) = \min\{p^* \mid p \in P, \varepsilon_p = 0\}$. If $\varepsilon \neq (0, \dots, 0)$, the father of I_ε in the positive greedy flip tree $\mathcal{P}(\mathcal{Q}^{\text{dup}})$ is the facet I_{ε^+} , where ε^+ is obtained from ε changing its last 1 into a 0. Similarly, if $\varepsilon \neq (1, \dots, 1)$, the father of I_ε in the negative greedy flip tree $\mathcal{N}(\mathcal{Q}^{\text{dup}})$ is the facet I_{ε^-} , where ε^- is obtained from ε changing its first 0 into a 1.

5.6. Minkowski sum decompositions into Coxeter matroid polytopes. Remember that the Minkowski sum of two polytopes $\Pi_1, \Pi_2 \subset V$ is the polytope

$$\Pi_1 + \Pi_2 := \{v_1 + v_2 \mid v_1 \in \Pi_1, v_2 \in \Pi_2\} \subset V.$$

A vertex v of $\Pi_1 + \Pi_2$ maximizing a linear functional $f : V \rightarrow \mathbb{R}$ on $\Pi_1 + \Pi_2$ is a sum $v_1 + v_2$ of vertices $v_1 \in \Pi_1$ and $v_2 \in \Pi_2$ maximizing f on Π_1 and Π_2 respectively.

Proposition 5.19. *The brick polytope $\mathcal{B}(\mathcal{Q})$ is the Minkowski sum of the polytopes*

$$\mathcal{B}(\mathcal{Q}, k) := \text{conv} \{w(I, k) \mid I \text{ facet of } \mathcal{S}(\mathcal{Q})\}$$

over all positions $k \in [m]$. In other words, the sum and convex hull operators in the definition of the brick polytope commute:

$$\mathcal{B}(\mathcal{Q}) := \text{conv}_I \sum_k w(I, k) = \sum_k \text{conv}_I w(I, k) =: \sum_k \mathcal{B}(\mathcal{Q}, k),$$

where the index k of the sums ranges over the positions $[m]$ and the index I of the convex hulls ranges over the facets of the subword complex $\mathcal{S}(\mathcal{Q})$.

Proof. The proof is identical to that of [PS12, Proposition 3.26]. We repeat it here for the convenience of the reader.

First, by definition of the brick vector, the brick polytope $\mathcal{B}(\mathbf{Q})$ is clearly a subset of the Minkowski sum $\sum_k \mathcal{B}(\mathbf{Q}, k)$. To prove equality, we only need to show that any vertex of $\sum_k \mathcal{B}(\mathbf{Q}, k)$ is also a vertex of $\mathcal{B}(\mathbf{Q})$.

Consider a linear functional $f : V \rightarrow \mathbb{R}$. For any two adjacent facets I, J of $\mathcal{S}(\mathbf{Q})$ with $I \setminus i = J \setminus j$, and any $k \in [m]$, we obtain that

- if $\mathbf{w}(I, k) \notin \{\pm \mathbf{w}(I, i)\}$, then $f(\mathbf{w}(I, k)) = f(\mathbf{w}(J, k))$;
- otherwise, $f(\mathbf{w}(I, k)) - f(\mathbf{w}(J, k))$ has the same sign as $f(\mathbf{B}(I)) - f(\mathbf{B}(J))$ by Lemma 4.5.

Consequently, a facet I_f maximizes $f(\mathbf{B}(\cdot))$ among all facets of $\mathcal{S}(\mathbf{Q})$ if and only if it maximizes $f(\mathbf{w}(\cdot, k))$ for any $k \in [m]$.

Now let v be any vertex of $\sum_k \mathcal{B}(\mathbf{Q}, k)$, and let $f_v : V \rightarrow \mathbb{R}$ denote a linear functional maximized by v on $\sum_k \mathcal{B}(\mathbf{Q}, k)$. Then v is the sum of the vertices in $\mathcal{B}(\mathbf{Q}, k)$ which maximize f_v . Thus, $v = \sum_k \mathbf{w}(I_f, k) = \mathbf{B}(I_f)$ is a vertex of $\mathcal{B}(\mathbf{Q})$. $\square$

Moreover, we observe that each summand $\mathcal{B}(\mathbf{Q}, k)$ is a W -matroid polytope as defined in [BGW97]. A *W-matroid polytope* is a polytope such that the group generated by the mirror symmetries of its edges is a subgroup of W . Equivalently, a W -polytope is a polytope whose edges are directed by the roots of Φ , and whose vertex set lies on a sphere centered at the origin. The edges of each summand $\mathcal{B}(\mathbf{Q}, k)$ are parallel to the roots since the edges of their Minkowski sum $\mathcal{B}(\mathbf{Q})$ are, and the vertices of $\mathcal{B}(\mathbf{Q}, k)$ are equidistant from the origin since they are all images of w_{q_k} under the orthogonal group W . Thus, each summand $\mathcal{B}(\mathbf{Q}, k)$ is indeed a W -matroid polytope.

Example 5.20. For the word $\mathbf{Q}^{\text{ex}}$ of Example 2.5, the Minkowski sum is formed by five points $\mathcal{B}(\mathbf{Q}^{\text{ex}}, k)$ for $k \in \{1, 2, 3, 5, 8\}$, three segments $\mathcal{B}(\mathbf{Q}^{\text{ex}}, 4) = \text{conv}\{e_2, e_4\}$, $\mathcal{B}(\mathbf{Q}^{\text{ex}}, 6) = \text{conv}\{\mathbb{1} - e_1, \mathbb{1} - e_3\}$ and $\mathcal{B}(\mathbf{Q}^{\text{ex}}, 9) = \text{conv}\{\mathbb{1} - e_3, \mathbb{1} - e_4\}$, and one triangle $\mathcal{B}(\mathbf{Q}^{\text{ex}}, 7) = \text{conv}\{e_1 + e_2, e_2 + e_3, e_2 + e_4\}$.

Example 5.21 (Duplicated word). Consider the word $\mathbf{Q}^{\text{dup}}$ defined in Example 3.8. According to our description of the weight function $\mathbf{w}(I_\varepsilon, \cdot)$ given in Example 4.3, we obtain that

- for $k \in [N]$, the polytope $\mathcal{B}(\mathbf{Q}^{\text{dup}}, k^*)$ is the single point ω_k ;
- for $p \in P$, the polytope $\mathcal{B}(\mathbf{Q}^{\text{dup}}, p^* + 1)$ is the segment from ω_p to $\omega_p + \alpha_p$.

Thus, the polytope $\mathcal{B}(\mathbf{Q}^{\text{dup}})$ is the Minkowski sum of $\Theta := \sum_{k \in [N]} \omega_k + \sum_{p \in P} \omega_p$ with the roots $\{\alpha_p \mid p \in P\}$ (considered as one dimensional polytopes), that is, a parallelepiped whose edges are the roots $\{\alpha_p \mid p \in P\}$.

6. CLUSTER COMPLEXES AND GENERALIZED ASSOCIAHEDRA REVISITED

The *cluster complex* as defined in [FZ03] encodes the exchange graph of clusters in cluster algebras. Its first polytopal realization was obtained in [CFZ02]. Later on, an independent realization was presented in [HLT11], based on the c -Cambrian fans [Rea06, Rea07b, RS09]. Recently S. Stella generalized in [Ste11] the approach of [CFZ02] and showed that the resulting realizations of the associahedra coincide with that of [HLT11].

In this section, we first study the brick polytope for the cluster complex as a spherical subword complex as defined in [CLS11]. In particular, we show that

the polar of the brick polytope indeed realizes the cluster complex, and provide a description of its normal vectors and facets. As a consequence, we show that the brick polytope and the realization in [HLT11] differ only by an affine translation. This opens new perspectives on the latter realization, and provides new proofs of several constructions therein, in particular their connections with the Cambrian lattices and fans studied in [Rea06, Rea07b, RS09, RS11].

6.1. Realizing cluster complexes. Let c be a *Coxeter element*, i.e. the product of all simple reflections in a given order. Following [Rea07b], we choose an arbitrary reduced expression c of c and let $w(c)$ denote the *c-sorting word* of w , i.e. the lexicographically first (as a sequence of positions) reduced subword of c^∞ for w . In particular, let $w_o(c)$ denote the c -sorting word of the longest element $w_o \in W$. It is proved in [CLS11] that the subword complex $\mathcal{S}(cw_o(c))$ is isomorphic to the cluster complex. Let $N := \ell(w_o) = |\Phi^+|$, and let $m := n + N$ be the length of $cw_o(c)$. The following theorem is now an easy consequence of the results developed in this paper.

Theorem 6.1. *The polar of the brick polytope $\mathcal{B}(cw_o(c))$ realizes the subword complex $\mathcal{S}(cw_o(c))$.*

Proof. The subset $[n]$ of positions of $[m]$ forms a facet of the $\mathcal{S}(cw_o(c))$ since its complement $w_o(c)$ is a reduced expression for w_o . Its root configuration $R([n])$ coincides with the root basis Δ . Thus, the word $cw_o(c)$ is realizing, and Theorem 4.8 ensures that the subword complex $\mathcal{S}(cw_o(c))$ is isomorphic to the boundary complex of the polar of the brick polytope $\mathcal{B}(cw_o(c))$. $\square$

Example 6.2. We illustrate the results of this section with the Coxeter element $c^{\text{ex}} := \tau_1\tau_2\tau_3$ of A_3 . The c^{ex} -sorting word of w_o is $w_o(c^{\text{ex}}) = \tau_1\tau_2\tau_3\tau_1\tau_2\tau_1$. We thus consider the word $c^{\text{ex}}w_o(c^{\text{ex}}) := \tau_1\tau_2\tau_3\tau_1\tau_2\tau_3\tau_1\tau_2\tau_1$. The facets of the subword complex $\mathcal{S}(c^{\text{ex}}w_o(c^{\text{ex}}))$ are $\{1, 2, 3\}$, $\{2, 3, 4\}$, $\{3, 4, 5\}$, $\{4, 5, 6\}$, $\{5, 6, 7\}$, $\{6, 7, 8\}$, $\{6, 8, 9\}$, $\{4, 6, 9\}$, $\{3, 5, 7\}$, $\{2, 4, 9\}$, $\{1, 3, 7\}$, $\{1, 7, 8\}$, $\{1, 8, 9\}$, and $\{1, 2, 9\}$. Their brick vectors and the brick polytope $\mathcal{B}(c^{\text{ex}}w_o(c^{\text{ex}}))$ are represented in Figure 12.




FIGURE 12. The brick polytope $\mathcal{B}(c^{\text{ex}}w_o(c^{\text{ex}}))$, for the Coxeter element $c^{\text{ex}} = \tau_1\tau_2\tau_3$, is the classical 3-dimensional associahedron.

6.2. Normal vectors and facets. We define the *weight configuration* of a facet I of $\mathcal{S}(\text{cw}_\circ(c))$ to be the set

$$W(I) := \{w(I, i) \mid i \in I\}$$

of all weights associated to the elements of I . Although not relevant for general words, the weight configuration is interesting for $\text{cw}_\circ(c)$ for the following reason.

Proposition 6.3. *For any facet I of $\mathcal{S}(\text{cw}_\circ(c))$, and any position $i \in I$, the weight $w(I, i)$ is a normal vector of the maximal face cone $\{r(I, j) \mid j \in I \setminus i\}$ of the cone $C(I)$. Thus, the normal cone $C^\circ(I)$ of $B(I)$ in $\mathcal{B}(\text{cw}_\circ(c))$ is generated by the weight configuration $W(I)$ of I .*

Proof. We derive this result from Proposition 3.9 and Lemma 5.2 in [CLS11]. The first proposition gives a rule to cyclically rotate the word $\text{cw}_\circ(c)$ until the letter i comes first, while maintaining the combinatorics of the subword complex. The second proposition ensures that, when $1 \in I$, the facet I contains only letters j for which $r(I, j)$ belongs to $\Phi_{\langle \alpha_{e_1} \rangle}$. Thus, $w(I, 1)$ is orthogonal to all the roots of $R(I) \setminus r(I, 1)$. $\square$

Let $w_1, \dots, w_N$ denote the letters of the c -sorting word of $w_\circ$. For $k \in [N]$, define $\rho_k := w_1 \cdots w_k \in W$ to be the product of the first k letters of $w_\circ(c)$ (with the convention $\rho_0 := e$). Moreover, let $\alpha_k := \rho_{k-1}(\alpha_{w_k})$, let $\omega_k := \rho_{k-1}(\omega_{w_k})$, and let $\Omega := \sum_{k \in [N]} \omega_k$. Remember that $\nabla := \{\omega_s \mid s \in S\}$ denotes the set of fundamental weights. The following observation is a consequence of Proposition 6.3.

Corollary 6.4. *The normal vectors of the facets of the brick polytope $\mathcal{B}(\text{cw}_\circ(c))$ are given by $\nabla \cup \{\omega_k \mid k \in [N]\}$.*

Proof. For any facet I of $\mathcal{S}(\text{cw}_\circ(c))$ and any $i \in I$, the corresponding weight $w(I, i)$ is by definition of the form ω_s , for some $s \in S$, or ω_k , for some $k \in [N]$. $\square$

To obtain the inequality description of the brick polytope $\mathcal{B}(\text{cw}_\circ(c))$, we now need to study carefully the facets $\kappa(\rho_k)$ of $\mathcal{S}(\text{cw}_\circ(c))$. The following lemma gives their root configurations and brick vectors.

Lemma 6.5. *For any $k \in \{0, \dots, N\}$, the root configuration and brick vector of $\kappa(\rho_k)$ are given by*

$$R(\kappa(\rho_k)) = \rho_k(\Delta) \quad \text{and} \quad B(\kappa(\rho_k)) = \Omega + \rho_k \left(\sum_{s \in S} \omega_s \right).$$

Proof. For this proof, we need some additional definitions. In the word $\text{cw}_\circ(c)$, we say that a position i is *terminal* if there is no position $j > i$ such that the i^{th} and j^{th} letters coincide. Let T denote the set of terminal positions. For $k \in [N]$, let $j_k = k + n + 1$, and let i_k denote the position of the last occurrence of the j_k^{th} letter before j_k . Note that $k + 1 \leq i_k \leq k + n$, that $i_k + 1 \leq i_{k+1}$, and that all positions i with $i_k + 1 \leq i < i_{k+1}$ are terminal.

We prove by induction on k that $\kappa(\rho_k) = (T \cap [i_k - 1]) \cup \{i_k, \dots, j_k - 1\}$ and that $r(\kappa(\rho_k), i_k) = r(\kappa(\rho_k), j_k) = \alpha_{k+1}$. The result is clear for $k = 0$, for which $\kappa(e) = [n]$, $i_0 = 1$ and $j_0 = n + 1$. Assume that it holds for a given $k \in [N]$. Since $\rho_{k+1} = \rho_k w_{(k+1)}$, we derive from Lemma 5.3 that

$$\begin{aligned} \kappa(\rho_{k+1}) &= (\kappa(\rho_k) \setminus i_k) \cup j_k = (T \cap [i_k - 1]) \cup \{i_k + 1, \dots, j_k\} \\ &= (T \cap [i_{k+1} - 1]) \cup \{i_{k+1}, \dots, j_{k+1} - 1\}. \end{aligned}$$

The inclusion $\{i_{k+1}, \dots, j_{p+1} - 1\} \subset \kappa(\rho_{k+1})$ ensures that $\sigma_{\kappa(\rho_{k+1})}^{[1, i_{p+1}-1]} = \sigma_{\kappa(\rho_{k+1})}^{[1, j_{p+1}-1]}$, and thus $r(\kappa(\rho_{k+1}), i_{k+1}) = r(\kappa(\rho_{k+1}), j_{k+1})$. The inclusion $\kappa(\rho_{k+1}) \subset [j_k]$ ensures that $\sigma_{\kappa(\rho_{k+1})}^{[1, j_k]} = \rho_{k+1}$, and thus $r(\kappa(\rho_{k+1}), j_{k+1}) = \rho_{k+1}(\alpha_{w_{(k+2)}}) = \alpha_{k+2}$. This proves that the result holds for $k+1$.

From our description of the facet $\kappa(\rho_k)$, we now derive its brick vector $B(\kappa(\rho_k))$. First, using that $\kappa(\rho_k) \setminus i_k = \kappa(\rho_{k+1}) \setminus j_k$, that $\{i_k, \dots, j_k - 1\} \subset \kappa(\rho_k)$, and that the i_k^{th} and j_k^{th} letters of $\text{cw}_o(c)$ coincide, we observe that the reduced expression of w_o given by the complement of $\kappa(\rho_k)$ is always $w_o(c)$. This ensures that $\sum_{k \notin \kappa(\rho_k)} w(\kappa(\rho_k), k) = \sum_{k' \in [N]} \omega_{k'} =: \Omega$. Assume now that $i \in \kappa(\rho_k)$, and let s denote the i^{th} letter of $\text{cw}_o(c)$. If $i \in \{i_k, \dots, j_k - 1\}$, then we have $w(\kappa(\rho_k), i) = \sigma_{\kappa(\rho_k)}^{[1, i_k-1]}(\omega_s) = \rho_k(\omega_s)$ since $\{i_k, \dots, j_k - 1\} \subset \kappa(\rho_k) \subset [j_k - 1]$. If i is terminal, then $w(\kappa(\rho_k), i) = w_o(\omega_s) = \rho_k(\omega_s)$. In any case, $w(\kappa(\rho_k), i) = \rho_k(\omega_s)$ for all $i \in \kappa(\rho_k)$. Since each simple reflection $s \in S$ appears in the word $\text{cw}_o(c)$ either at a position in $\{i_k, \dots, j_k - 1\}$ or at a terminal position before i_k , we obtain that $W(\kappa(\rho_k)) = \rho_k(\nabla)$, and thus $\sum_{i \in \kappa(\rho_k)} w(\kappa(\rho_k), i) = \rho_k(\sum_{s \in S} \omega_s)$. Summing the two contributions, we derive the brick vector

$$B(\kappa(\rho_k)) := \sum_{k \in [m]} w(\kappa(\rho_k), k) = \Omega + \rho_k \left(\sum_{s \in S} \omega_s \right).$$

Finally, according to Lemma 4.5, the root configuration $R(\kappa(\rho_k))$ is the set of roots orthogonal to the facet defining hyperplanes of the normal cone of $B(\kappa(\rho_k))$ in $\mathcal{B}(\text{cw}_o(c))$. Since this cone is generated by $W(\kappa(\rho_k)) = \rho_k(\nabla)$, we obtain that $R(\kappa(\rho_k)) = \rho_k(\Delta)$. $\square$

Let $q := \sum_{s \in S} \omega_s$ and let $\mathcal{P}(W) := \text{conv} \{w(q) \mid w \in W\}$ be the balanced W -permutahedron.

Theorem 6.6. *The brick polytope $\mathcal{B}(\text{cw}_o(c))$ is obtained from the balanced permutahedron $\mathcal{P}(W)$ removing all facets which do not intersect $\{\rho_k(q) \mid 0 \leq k \leq N\}$, and then translating by the vector $\Omega := \sum_{k \in [N]} \omega_k$.*

Proof. Let $\mathcal{B}'$ denote the translate of $\mathcal{B}(\text{cw}_o(c))$ by the vector $-\Omega$. According to Lemma 6.5, the vertices of $\mathcal{B}'$ and $\mathcal{P}(W)$ corresponding to the prefixes ρ_k of $w_o(c)$ coincide, as well as their normal cones. Thus, the facets of $\mathcal{B}'$ and $\mathcal{P}(W)$ containing these vertices are defined by the same inequalities. The result thus follows from Corollary 6.4 which affirms that these inequalities exhaust all facets of $\mathcal{B}'$. $\square$

Corollary 6.7. *Up to the translation by the vector Ω , the brick polytope $\mathcal{B}(\text{cw}_o(c))$ coincides with the c -associahedron $\text{Asso}_c^q(W)$ in [HLT11].*

Remark 6.8. By a recent work of S. Stella [Ste11], the c -associahedron $\text{Asso}_c^q(W)$ also coincides, up to an affine translation, with the first realization of the cluster complex by F. Chapoton, S. Fomin, and A. Zelevinsky [CFZ02] and its generalization by S. Stella [Ste11]. We thus obtained in the present paper the vertex description of both realizations (by Definition 4.1) and a natural Minkowski sum decomposition into W -matroid polytopes (by Proposition 5.19).

Example 6.9. For the Coxeter element $c^{\text{ex}} := \tau_1 \tau_2 \tau_3$ of Example 6.2, the translation vector is $\Omega = (3, 2, 3, 6)$. We have represented in Figure 13 the balanced 3-dimensional permutahedron $\mathcal{P}(A_3) := \text{conv} \{\sigma(0, 1, 2, 3) \mid \sigma \in \mathfrak{S}_4\}$ together with




FIGURE 13. The brick polytope $\mathcal{S}(c^{\text{ex}}w_o(c^{\text{ex}}))$ is obtained removing facets of the balanced permutahedron $\mathcal{P}(A_3)$. The remaining facets of the permutahedron are shaded.

the 3-dimensional associahedron $\mathcal{B}(c^{\text{ex}}w_o(c^{\text{ex}})) - \Omega$. The set $\{\rho_k(q) \mid 0 \leq k \leq 6\}$ are all the vertices located on the leftmost path from the bottommost vertex to the topmost vertex in the graphs of these polytopes (they are marked with strong red dots). The facets of the permutahedron $\mathcal{P}(A_3)$ containing these points, which give precisely the facets of the associahedron $\mathcal{B}(c^{\text{ex}}w_o(c^{\text{ex}})) - \Omega$, are shaded.

In the end of this paper, we revisit various results contained in [HLT11]. The approach developed in this paper, based on subword complexes and brick polytopes, enables us to simplify certain proofs and connections.

First, we provide a new way to see that the c -singletons as defined in [HLT11] can be used to characterize the vertices common to the brick polytope $\mathcal{B}(cw_o(c))$ and to the balanced W -permutahedron $\mathcal{P}(W)$. In the following list, immediate reformulations of the same characterization are indicated by a — symbol.

Proposition 6.10. *The following properties are equivalent for an element $w \in W$:*

- (i) *The element w is a **singleton** for the map κ , i.e. $\kappa^{-1}(\kappa(w)) = \{w\}$.*
- (ii) *The root configuration of $\kappa(w)$ is given by $R(\kappa(w)) = w(\Delta)$.*
 - *The root configuration $R(\kappa(w))$ of $\kappa(w)$ is a **simple system** for W , that is, the set of extremal roots of a separable subset of Φ .*
 - *The cone $C(\kappa(w))$ of $\mathcal{B}(cw_o(c))$ at $B(\kappa(w))$ is the polar cone of the chamber $w(\mathcal{C})$ of the Coxeter arrangement of W , i.e. $C(\kappa(w)) = w(\mathcal{C})^\circ = w(\mathcal{C}^\circ)$.*
- (iii) *The weight configuration $W(\kappa(w))$ of $\kappa(w)$ is given by $W(\kappa(w)) = w(\nabla)$.*
 - *The weight configuration $W(\kappa(w))$ of $\kappa(w)$ generates a chamber of the Coxeter arrangement of W .*
 - *The normal cone $C^\circ(\kappa(w))$ of $B(\kappa(w)) \in \mathcal{B}(cw_o(c))$ coincides with the chamber $w(\mathcal{C})$ of the Coxeter arrangement of W , i.e. $C^\circ(\kappa(w)) = w(\mathcal{C})$.*
- (iv) *The vertices $B(\kappa(w))$ of the brick polytope $\mathcal{B}(cw_o(c))$ and $w(q)$ of the balanced W -permutahedron $\mathcal{P}(W)$ coincide up to Ω , i.e. $B(\kappa(w)) = \Omega + w(q)$.*
- (v) *There exist reduced expressions w of w and c of c such that w is a prefix of $w_o(c)$.*
- (vi) *The complement of $\kappa(w)$ in $cw_o(c)$ is $w_o(c)$.*




FIGURE 14. The c -associahedron of type A_3 for the Coxeter element c being $\tau_1\tau_2\tau_3$ (top left), $\tau_3\tau_2\tau_1$ (top right), $\tau_2\tau_1\tau_3$ (bottom left), and $\tau_3\tau_1\tau_2$ (bottom right).





FIGURE 15. The c -associahedron of type B_3 for the Coxeter element c being $\tau_2\tau_1\chi$ (top) and $\tau_2\chi\tau_1$ (bottom).

FIGURE 16. An H_3 -associahedron.

Proof. Properties (i), (ii), and (iii), as well as their respective variants, are equivalent according to Lemma 5.3, Proposition 5.4, and Proposition 6.3.

If w violates (v), then at least one of the weights of $w(\nabla)$ does not belong to $\nabla \cup \{\omega_k \mid k \in [N]\}$. Proposition 6.3 then affirms that w also violates (iii). Thus, (iii) implies (v).

We proved in Lemma 6.5 that any prefix of $w_\circ(c)$ satisfies (iv). Any two reduced expressions c and c' for c are related by a sequence of commutations of letters s, t with $m_{st} = 1$. It induces an isomorphism between the subword complexes $\mathcal{S}(cw_\circ(c))$ and $\mathcal{S}(c'w_\circ(c'))$, which commutes with the root configuration $R(\cdot)$, the brick vector $B(\cdot)$ and the map κ . In particular, the functions $R(\kappa(\cdot))$ and $B(\kappa(\cdot))$ are independent of the choice of the reduced expression c of c . Consequently, Lemma 6.5 ensures that (v) implies (iv).

Similarly, we have seen in the proof of Lemma 6.5 that (vi) holds when w is a prefix of $w_\circ(c)$. Up to commutations, we thus have that (v) implies (vi).

Since the permutahedron $\mathcal{P}(W)$ is a simple polytope, the only vertices which can survive when removing a set $\mathcal{F}$ of facets of $\mathcal{P}(W)$ are those which do not belong to any facet of $\mathcal{F}$. Consequently, Theorem 6.6 ensures that (iv) implies (iii).

Finally, if the complement of $\kappa(w)$ in $cw_\circ(c)$ is precisely $w_\circ(c)$, then the weights of $\kappa(w)$ all lie in $\nabla \cup \{\omega_k \mid k \in [N]\}$. Thus, all facets incident to $\kappa(w)$ when deleting the facets of the permutahedron, and (vi) implies (iii). $\square$

Remark 6.11. For completeness, we want to point out that all definitions and results of Sections 3 and 4 easily generalize if we replace $r(I, k)$ by $\lambda(q_k)r(I, k)$ and $w(I, k)$ by $\lambda(q_k)w(I, k)$ for an arbitrary function $\lambda : S \rightarrow \mathbb{R}_{>0}$. The resulting polytope $\mathcal{B}_\lambda(Q)$ has the same combinatorial properties for any λ , but its geometry is different. We decided to present our construction and results with $\lambda \equiv 1$ to simplify the exposition. In the special case of the subword complex $\mathcal{S}cw_\circ(c)$, the resulting brick polytope $\mathcal{B}_\lambda(cw_\circ(c))$ is a translate of the c -associahedron $\text{Asso}_c^x(W)$ in [HLT11], where $x = \sum_{s \in S} \lambda(s)\omega_s$. This polytope is obtained from the permutahedron $\mathcal{P}_x(W)$ removing all facets which do not intersect $\{\rho_k(x) \mid 0 \leq k \leq N\}$.

Example 6.12. Figures 14, 15, and 16 represent type A_3 , B_3 , and H_3 associahedra obtained by removing facets of the type A_3 , B_3 , and H_3 permutahedra of Figures 2,

3, and 4. They are all obtained from the corresponding permutahedron removing facets not containing c -singletons, which are marked with dots.

Remark 6.13. In regard of the results of this section, it is tempting to try to generalize the geometric description of the generalized associahedra to all brick polytopes. We observe however that there exist realizing words Q for which

- the weight configuration $W(I) = \{w(I, i) \mid i \in I\}$ of a facet I of $\mathcal{S}(Q)$ does not necessarily generate the normal cone of $B(I)$ in $\mathcal{B}(Q)$.
- no element $w \in W$ is a singleton for the map κ . In other words, no chamber of the Coxeter fan is the normal cone of a vertex of the brick polytope $\mathcal{B}(Q)$. In particular, the chambers $\mathcal{C}$ and $w_o(\mathcal{C})$ are not necessarily normal cones of the brick polytope.
- no translate of the brick polytope $\mathcal{B}(Q)$ is obtained from a W -permutahedron by deleting facets.

6.3. Five Coxeter-Catalan families. In this section, we temporarily interrupt our geometric study of the cluster complex to describe and relate the combinatorics of several Coxeter-Catalan families. We will need these relations in the following section to connect brick polytopes to Cambrian lattices and fans. We revisit here the bijective connections between

- (i) the set $\text{CLUS}_c(W)$ of c -clusters,
- (ii) the set $\text{SORT}_c(W)$ of c -sortable elements,
- (iii) the set $\text{NCP}_c(W)$ of c -noncrossing partitions,
- (iv) the set $\text{NCS}_c(W)$ of c -noncrossing subspaces,

and describe their connection to the set of facets of $\mathcal{S}(\text{cw}_o(c))$ which we denote for convenience in this section by $\text{FAC}_c(W)$. The bijective connections are summarized in Figure 17.




FIGURE 17. Coxeter-Catalan families and their connections.

All connections between these Coxeter-Catalan families are based on their analogous inductive structure, described below. Although the proofs really rely on this inductive structure, most maps between also be described directly, avoiding the use of induction.

6.3.1. Facets of the subword complex and clusters. Since this connection is the most straightforward, we start with the identification between c -clusters and facets of $\mathcal{S}(\text{cw}_o(c))$ as given in [CLS11]. To this end, let $\Phi_{\geq -1} := -\Delta \cup \Phi^+$ be the set of *almost positive roots*. Denote by $w(c) := w_1 \cdots w_{\ell(w)}$ the *c -sorting word* of some

element $w \in W$, i.e. the lexicographically first reduced subword of c^∞ for w , and let $m := n + \ell(w)$ be the length of the word $\text{cw}(c)$. Associate to each position $i \in [m]$ in this word an almost positive root $\text{lr}_c(w, i)$ by

$$\text{lr}_c(w, i) := \begin{cases} -\alpha_{c_i} & \text{if } i \leq n, \\ w_1 \cdots w_{i-n-1}(\alpha_{w_{i-n}}) & \text{otherwise.} \end{cases}$$

In the case where w is the longest element $w_\circ$, it is shown in [CLS11, Theorem 2.2] that the map $\text{lr}_c(\cdot) := \text{lr}_c(w_\circ, \cdot)$ induces a bijection $\text{lr}_c : \text{FAC}_c(W) \rightarrow \text{CLUS}_c(W)$ between facets of $\mathcal{S}(\text{cw}_\circ(c))$ and c -clusters,

$$\text{lr}_c : \begin{array}{ccc} \text{FAC}_c(W) & \longrightarrow & \text{CLUS}_c(W) \\ I & \longmapsto & \{\text{lr}_c(i) \mid i \in I\} \end{array}.$$

In other words, write once and for all the set of almost positive roots $\Phi_{\geq -1}$ as the list $\text{lr}_c(1), \dots, \text{lr}_c(m)$. Then the cluster $\text{lr}_c(I)$ associated to a facet I in $\text{FAC}_c(W)$ is given by its sublist at positions in I , while the facet $\text{lr}_c^{-1}(X)$ associated to a c -cluster X is the set of positions of X within this list. Note in particular that the elements in the c -cluster X are naturally ordered by their position in this list. We here use this description of c -clusters as the definition of the set $\text{CLUS}_c(W)$ and refer the reader to [Rea07a] and [CLS11] for further background.

Example 6.14. Consider the Coxeter element $c^{\text{ex}} := \tau_1 \tau_2 \tau_3$ of A_3 already discussed in Example 6.2. The c^{ex} -sorting word for $w_\circ$ is $c^{\text{ex}} w_\circ(c^{\text{ex}}) = \tau_1 \tau_2 \tau_3 \tau_1 \tau_2 \tau_1$. Therefore, the list $\text{lr}_{c^{\text{ex}}}(1), \dots, \text{lr}_{c^{\text{ex}}}(m)$ is given by

$$e_1 - e_2, e_2 - e_3, e_3 - e_4, e_2 - e_1, e_3 - e_1, e_4 - e_1, e_3 - e_2, e_4 - e_2, e_4 - e_3.$$

Consider now the facet $F^{\text{ex}} := \{4, 6, 9\} \in \text{FAC}_{c^{\text{ex}}}(A_3)$. Its associated cluster is given by $\text{lr}_{c^{\text{ex}}}(F^{\text{ex}}) = \{e_2 - e_1, e_4 - e_1, e_4 - e_3\}$, while its root configuration is given by $R(F^{\text{ex}}) = \{e_3 - e_2, e_1 - e_3, e_3 - e_4\}$. The following proposition gives a direct relation between these two sets of roots.

The next proposition connects the root configuration $R(I)$ to the c -cluster $\text{lr}_c(I)$ associated to a facet $I \in \text{FAC}_c(W)$. This connection enables us to compute $R(I)$ from $\text{lr}_c(I)$ and vice versa. We will need it later in the description of the bijection $\text{rp}_c : \text{CLUS}_c(W) \rightarrow \text{NCP}_c(W)$.

Proposition 6.15. *Let $I := \{i_1 < \dots < i_n\}$ be a facet in $\text{FAC}_c(W)$. Consider its root configuration $R(I) = \{\alpha_1, \dots, \alpha_n\}$, where $\alpha_p := r(I, i_p)$, and the associated c -cluster $\text{lr}_c(I) = \{\beta_1, \dots, \beta_n\}$, where $\beta_p := \text{lr}_c(i_p)$. Moreover, let j be such that $i_j \leq n$ and $i_{j+1} > n$, or in other words, $\{\beta_1, \dots, \beta_j\} \subseteq -\Delta$, $\{\beta_{j+1}, \dots, \beta_n\} \subseteq \Phi_{\langle \beta_1, \dots, \beta_j \rangle}^+$. For any p with $j < p \leq n$, we have*

$$(1) \quad \alpha_p = -s_{\beta_n} \cdots s_{\beta_{p+1}}(\beta_p) \quad \text{and} \quad \beta_p = -s_{\alpha_n} \cdots s_{\alpha_{p+1}}(\alpha_p).$$

Moreover, we have

$$(2) \quad s_{\alpha_{j+1}} \cdots s_{\alpha_n} = s_{\beta_n} \cdots s_{\beta_{j+1}} = c',$$

where c' is obtained from c by removing all letters $s_{\beta_1}, \dots, s_{\beta_j}$.

Proof. Starting with $\alpha_n = -\beta_n$, it is straightforward to check that both statements in (1) are equivalent. We thus only show the second equality. We write

$\text{cw}_\circ(c) := q_1 \cdots q_m$, and observe that by definition,

$$(*) \quad \begin{aligned} \beta_p &= w_\circ q_m q_{m-1} \cdots q_{i_p}(\alpha_{q_{i_p}}), \\ \alpha_p &= w_\circ q_m q_{m-1} \cdots \hat{q}_{i_n} \cdots \hat{q}_{i_{p+1}} \cdots q_{i_p+1}(\alpha_{q_{i_p}}), \end{aligned}$$

where the hats on the $\hat{q}_{i_n}, \dots, \hat{q}_{i_{p+1}}$ mean that they are omitted in the expression for α_p . Thus applying successively $s_{\beta_n}, \dots, s_{\beta_{p+1}}$ to α_p inserts successively $q_{i_n}, \dots, q_{i_{p+1}}$ into the righthand side of $(*)$, statement (1) follows. To prove (2), observe that the first equality follows from (1) with $p = j + 1$. The second equality is directly deduced from [CLS11, Proposition 2.8] with $k = 1$. $\square$

6.3.2. Facets of the subword complex and sortable elements. Recall the definition of c -sortable elements from [Rea07b]. Let w be an element of W . Its c -sorting word can be written as $w(c) = c_{K_1} c_{K_2} \cdots c_{K_p}$, where c_K denotes the subword of c only taking the simple reflections in $K \subset S$ into account. The element w is called *c -sortable* if its c -sorting word $w(c) = c_{K_1} c_{K_2} \cdots c_{K_p}$ is nested, *i.e.* if $K_1 \supseteq K_2 \supseteq \cdots \supseteq K_p$. Observe that the property of being c -sortable does not depend on the particular reduced expression c but only on c . We denote by $\text{SORT}_c(W)$ the set of c -sortable elements in W .

N. Reading defines in [Rea07b, Theorem 8.1] the bijection $\text{cl}_c : \text{SORT}_c(W) \longrightarrow \text{CLUS}_c(W)$ between c -sortable elements and c -clusters by

$$\text{cl}_c(w) := \{l_c(i) \mid w_i \text{ is the last occurrence of a letter in } \text{cw}(c)\}.$$

He also provides the following inductive description of c -sortable elements which is immediate from the definition. Let $w \in W$, and let s be *initial* in c , *i.e.* $\ell(sc) < \ell(c)$. Then

$$w \in \text{SORT}_c(W) \iff \begin{cases} sw \in \text{SORT}_{scs}(W) & \text{if } \ell(sw) < \ell(w), \\ w \in \text{SORT}_{sc}(W_{(s)}) & \text{if } \ell(sw) > \ell(w). \end{cases}$$

The first case yields an induction on the length $\ell(w)$, while the second case is an induction on the rank of W .

We now describe the analogue inductive nature of the facets of $\mathcal{S}(\text{cw}_\circ(c))$. To this end, we define the *rotated word* of $Q := q_1 q_2 \cdots q_m$ to be $Q_\circ := q_2 \cdots q_m \psi(q_1)$, where ψ denotes the automorphism on S given by $\psi(s) := w_\circ^{-1} s w_\circ$. Let s be initial in c , let c be a word for c starting with s , and let $I \in \text{FAC}_c(W)$. Then there are two cases.

- (i) The first letter s in $\text{cw}_\circ(c)$ is not contained in I . We then have by [CLS11, Proposition 4.3] that I , shifted by one, is a facet of the subword complex for the word $(\text{cw}_\circ(c))_\circ$, which is, up to commutations of consecutive commuting letters, equal to $c' w_\circ(c')$ with $c' := scs$.
- (ii) The first letter s in $\text{cw}_\circ(c)$ is contained in I . [CLS11, Lemma 5.2] then ensures that $k \notin I$ for all k such that $r(I, k) \in \Phi \setminus \Phi_{\langle \alpha_s \rangle}$. By [CLS11, Lemma 5.4] (see also our Proposition 3.7), we therefore obtain that I can also be considered as a facet of the subword complex for the word $c' w_\circ(c')$ with $c' := sc$.

This analogous inductive structures yield a bijection

$$\text{so}_c := \text{cl}_c^{-1} \circ l_c : \text{FAC}_c(W) \longrightarrow \text{SORT}_c(W).$$

We have already seen two ways of computing so_c .

- (1) The first uses the inductive description: the c -sorting word w of $w = \text{so}_c(I)$ is obtained by looking at the first letter s of $\text{cw}_o(c)$. Depending on situations (i) and (ii) above, this letter is appended or not to w . Then the first letter is rotated, or all positions k for which $r(I, k) \in \Phi \setminus \Phi_{\langle \alpha_s \rangle}$ are deleted.
- (2) The second uses the iterative procedure described in Remark 5.12 and Proposition 5.13.

Observe that the first follows the same lines as the description in [Rea07b, Remark 6.9], see also Section 6.3.3 below.

Example 6.16. Consider the facet $F^{\text{ex}} := \{4, 6, 9\} \in \text{FAC}_{c^{\text{ex}}}(A_3)$ already mentioned in Example 6.14. Its root configuration is $R(F^{\text{ex}}) = \{e_3 - e_2, e_1 - e_3, e_3 - e_4\}$ so that $R^+(F^{\text{ex}}) = \{e_3 - e_2\}$ and $R^-(F^{\text{ex}}) = \{e_3 - e_1, e_4 - e_3\}$. Consequently, we have $\Lambda(F^{\text{ex}}) = \{e_3 - e_1, e_4 - e_3, e_4 - e_1, e_2 - e_1\}$, which is the inversion set of the c^{ex} -sortable element $\text{so}_c(F^{\text{ex}}) = 2431 = \tau_1\tau_2\tau_3\tau_2$. Figure 18 represents all facets of $\text{FAC}_{c^{\text{ex}}}(A_3)$ and their corresponding c^{ex} -sortable elements by the map $\text{so}_{c^{\text{ex}}}$.

We can now use the bijection so_c to transport structures from facets of $\mathcal{S}(\text{cw}_o(c))$ to c -sortable elements of W or vice versa.

We start with the canonical tree structures on these objects. As pointed out by N. Reading in [Rea07a, Remark 2.1], the c -sortable elements of W have a natural search-tree structure. The root of this tree is the identity element $e \in W$ and the father of a c -sortable element w is the c -sortable element whose c -sorting word is obtained from the c -sorting word of w deleting its last letter. We denote this tree by $\mathcal{T}(c)$ and call it the c -sorting tree. Note that the c -sorting tree depends on the particular reduced expression c for c , although the c -sortable elements do not. The following proposition, illustrated in Figure 18, connects the greedy flip tree defined in Section 5.5 with the c -sorting tree.

Proposition 6.17. *The bijection so_c is an isomorphism between the positive greedy flip tree $\mathcal{P}(\text{cw}_o(c))$ and the c -sorting tree $\mathcal{T}(c)$.*

Proof. The statement follows from the analogous inductive structures of these trees. $\square$

Similarly, the bijection so_c is an isomorphism between the negative greedy flip tree $\mathcal{P}(\text{cw}_o(c))$ and the c -antisortable tree.

Example 6.18. Figure 18 represents the c^{ex} -sorting tree $\mathcal{T}(c^{\text{ex}})$ and the positive greedy flip tree $\mathcal{P}(c^{\text{ex}}\text{w}_o(c^{\text{ex}}))$, for the Coxeter element $c^{\text{ex}} := \tau_1\tau_2\tau_3$ of Example 6.2. See also Figure 19 for a more geometric picture of the greedy flip trees of $\mathcal{S}(c^{\text{ex}}\text{w}_o(c^{\text{ex}}))$.

In the following, we use the inductive structure of $\mathcal{S}(\text{w}_o(c))$ to see that the root configuration was already present in [RS11]. Therein, N. Reading and D. Speyer defined inductively a set $\mathcal{C}_c(w)$ associated to a c -sortable element w ,

$$\mathcal{C}_c(w) := \begin{cases} s\mathcal{C}_{scs}(sw) & \text{if } \ell(sw) < \ell(w), \\ \mathcal{C}_{sc}(w) \cup \{\alpha_s\} & \text{if } \ell(sw) > \ell(w). \end{cases}$$

This set is intimately related to the root configuration.

Proposition 6.19. *For any facet $I \in \text{FAC}_c(W)$, the root configuration $R(I)$ and the set $\mathcal{C}_c(\text{so}_c(I))$ coincide.*




FIGURE 18. The bijection $\text{so}_{c^{\text{ex}}}$ sends the c^{ex} -sorting tree $\mathcal{T}(c^{\text{ex}})$ to the positive greedy flip tree $\mathcal{P}(c^{\text{ex}}w_o(c^{\text{ex}}))$. Each box corresponds to a facet I of $\mathcal{S}(c^{\text{ex}}w_o(c^{\text{ex}}))$: the bottom frame contains the facet I and the top frame contains the c^{ex} -sorting word of $\text{so}_{c^{\text{ex}}}(I)$.




FIGURE 19. The positive (left) and negative (right) greedy flip trees of the subword complex $\mathcal{S}(c^{\text{ex}}w_o(c^{\text{ex}}))$ of Example 2.5, seen as monotone spanning trees of the 1-skeleton of the associahedron $\mathcal{B}(c^{\text{ex}}w_o(c^{\text{ex}}))$.

Proof. This is a direct consequence of the inductive description of c -sortable elements and of the facets of $\mathcal{S}(\text{cw}_o(c))$ presented above. $\square$

Proposition 6.19 yields the following description of $\mathcal{C}_c(w)$. Let $w \in W$ be c -sortable with c -sorting word $w := w_1 \cdots w_{\ell(w)}$, let $s \in S$, and let $k \in [\ell(w)]$. We say that w *skips* s in position $k+1$ if the leftmost instance of s in c^∞ not used in w occurs after position k . A skip s at position $k+1$ is moreover called *forced* if $w_1 \cdots w_k(\alpha_s) \in \Phi^-$, and *unforced* otherwise.

Corollary 6.20 ([RS11, Proposition 5.1]). *We have*

$$\mathcal{C}_c(w) = \{w_1 \cdots w_k(\alpha_s) \mid w \text{ skips } s \text{ at position } k+1\}$$

and the forced and unforced skips are given by the intersection of $\mathcal{C}_c(w)$ with Φ^- and with Φ^+ , respectively.

Remark 6.21. Remark 3.5 yields a shortcut for $\text{so}_c^{-1} = \text{lr}_c^{-1} \circ \text{cl}_c : \text{SORT}_c(W) \rightarrow \text{FAC}_c(W)$ using Proposition 6.19 and its Corollary 6.20. Observe moreover that the distinction between forced and unforced skips naturally arises when we sweep the word $\text{cw}_o(c)$ from left to right as described in Remark 3.5.

6.3.3. Facets of the subword complex and noncrossing partitions. We start with recalling the definition of noncrossing partitions as given for example in [Rea07b, Section 5] (see also Lemma 5.2 therein for an inductive description). Define the *reflection length* $\ell_R(w)$ for $w \in W$ to be the length of the shortest factorization of w into reflections. Observe that in particular, the reflection length of any Coxeter element c is n . An element $w \in W$ is a *noncrossing partition* if $\ell_R(w) + \ell_R(w^{-1}c) = n$. The set of all noncrossing partitions is denoted by $\text{NCP}_c(W)$. The term *partition* has historical reasons and comes from the connection to noncrossing set partitions in the classical types. The c -sorting word $w(c) := w_1 \cdots w_{\ell(w)}$ of some element $w \in W$ induces a total order on its inversion set

$$\text{inv}(w) := \{\alpha_{w_1}, w_1(\alpha_{w_2}), \dots, w_1 w_2 \cdots w_{\ell(w)-1}(\alpha_{w_{\ell(w)}})\} \subset \Phi^+.$$

The (ordered) set of *cover reflections* of w is given by

$$\text{cov}(w) := \{t \in R \mid tw = ws < w \text{ for some } s \in S\} \subseteq \{s_\beta \mid \beta \in \text{inv}(w)\}.$$

In [Rea07b, Theorem 6.1], N. Reading showed that the map

$$\text{nc}_c : \begin{array}{ccc} \text{SORT}_c(W) & \longrightarrow & \text{NCP}_c(W) \\ w & \longmapsto & t_1 \cdots t_k \end{array}$$

is a bijection, where $\{t_1, \dots, t_k\} = \text{cov}(w)$ is the ordered set of cover reflections of w .

Using this result and the description of $\mathcal{C}_c(w)$ in the previous section, we are now in the situation to provide a direct connection between facets in $\text{FAC}_c(W)$ and noncrossing partitions. A position i in a facet $I \in \text{FAC}_c(W)$ is an *upper position* in I if $r(I, i) \in \Phi^-$. Moreover, its corresponding almost positive root $\text{lr}_c(i)$ is an *upper root* in $\text{lr}_c(I)$. Compare the original equivalent definition of upper roots before Corollary 8.6 in [RS09].

Theorem 6.22. *The bijection $\text{lp}_c = \text{nc}_c \circ \text{so}_c$ is given by*

$$\text{lp}_c : \begin{array}{ccc} \text{FAC}_c(W) & \longrightarrow & \text{NCP}_c(W) \\ I & \longmapsto & s_{r(I, j_1)} \cdots s_{r(I, j_k)}, \end{array}$$

where $\{j_1 < \cdots < j_k\}$ is the collection of upper positions in I .

Proof. The construction of $\text{so}_c : \text{FAC}_c(W) \rightarrow \text{SORT}_c(W)$ implies that the sorted set of cover reflections of $\text{so}_c(I)$ is given by $R(I) \cap \Phi^-$. This can as well be deduced from [RS11, Proposition 5.2] together with our Proposition 6.19. The statement then follows from the description of the map $\text{nc}_c : \text{SORT}_c(W) \rightarrow \text{NCP}_c(W)$ given above. $\square$

Example 6.23. Consider the facet $F^{\text{ex}} := \{4, 6, 9\} \in \text{FAC}_{c^{\text{ex}}}(A_3)$ mentioned in Examples 6.14 and 6.16. Its root configuration is $R(F^{\text{ex}}) = \{e_3 - e_2, e_1 - e_3, e_3 - e_4\}$. Consequently, its upper positions are 6 and 9 and its associated c -noncrossing partition is $\text{lp}_{c^{\text{ex}}}(F^{\text{ex}}) = (13)(34) = 3241$.

Theorem 6.24. *The bijection $\text{rp}_c = \text{lp}_c \circ \text{lr}_c^{-1}$ is given by*

$$\begin{aligned} \text{rp}_c : \text{CLUS}_c(W) &\longrightarrow \text{NCP}_c(W) \\ \{\beta_1, \dots, \beta_n\} &\longmapsto s_{\beta_{j_k}} \cdots s_{\beta_{j_1}}, \end{aligned}$$

where the product is taken over all upper roots from right to left (recall here that a c -cluster is naturally ordered, see Section 6.3.1).

Remark 6.25. Observe that Proposition 6.15 immediately implies that the bijection rp_c as given in Theorem 6.24 is well-defined.

Proof of Theorem 6.24. Let $I = \{i_1 < \cdots < i_n\}$ be the facet in $\text{FAC}_c(W)$ such that $\text{lr}_c(I)$ is given by $\{\beta_1, \dots, \beta_n\}$. The set of upper positions in I is then given by $\{j_1 < \cdots < j_k\} \subseteq I$. Observe moreover that in the situation of Proposition 6.15, all upper positions appear within $\{i_{j+1}, \dots, i_n\}$, and that the product $s_{\alpha_{j+1}} \cdots s_{\alpha_n}$ on the left-hand side of Proposition 6.15(2) can be reordered by commutations such that all upper positions appear last in this expression, i.e., that the product $s_{\alpha_{j_1}} \cdots s_{\alpha_{j_k}}$ appear as a suffix of $s_{\alpha_{j+1}} \cdots s_{\alpha_n}$. To see the later, it is enough to consider facets whose intersection with $[n]$ is empty. For these, it is straightforward to check that the upper positions appear after the others up to commutations. Together with Proposition 6.15(1), we thus obtain that

$$s_{\beta_{i_k}} \cdots s_{\beta_{i_1}} = s_{\text{lr}_c(i_k)} \cdots s_{\text{lr}_c(i_1)} = s_{r(I, i_1)} \cdots s_{r(I, i_k)}.$$

The statement follows. $\square$

Example 6.26. Consider the c^{ex} -cluster $X^{\text{ex}} := \{e_2 - e_1, e_4 - e_1, e_4 - e_3\}$. Its upper roots are $e_4 - e_1$ and $e_4 - e_3$. Consequently, its associated noncrossing partition is $\text{rp}_c(X^{\text{ex}}) = (34)(14) = 3241$.

Remark 6.27. The composition of rp_c with the map $\text{fix} : \text{NCP}_c(W) \rightarrow \text{NCS}_c(W)$ was described in [RS09, Theorem 11.1] using some further recursive notions, see also the paragraph thereafter. We here provide a direct map, not using the recursive structure. Theorem 6.24 provides as well new proofs of [RS09, Conjectures 11.2 and 11.3] which were as well proven in [IT09, Theorem 4.10].

Finally, we sketch an inductive description of $\text{sc}_c : \text{NCS}_c(W) \rightarrow \text{FAC}_c(W)$. To construct the image $\text{sc}_c(L)$ of a c -noncrossing subspace $L \in \text{NCS}_c(W)$, consider two cases depending on the last letter q_m of $\text{cw}_c(c)$. We set $s := \psi(q_m)$, which we can assume to be final in c , and we define $c' := scs$ and $c'' := cs = sc'$. The two cases depend on the inductive description of the facets describing how to embed $\text{FAC}_{c'}(W)$ and $\text{FAC}_{c''}(W_{\langle s \rangle})$ into $\text{FAC}_c(W)$.

- (i) If L is not contained in the reflection hyperplane of s , $\text{sc}_c(L)$ is given by the facet $\text{sc}_{c'}(sL)$ inside $\text{FAC}_{c'}(W)$.

- (ii) If L is contained in the reflection hyperplane of s , we obtain $\text{sc}_c(L)$ by adding the position m to the facet $\text{sc}_{c''}(L)$ inside $\text{FAC}_{c''}(W_{\langle s \rangle})$.

Observe here, that this procedure considers a final letter in c while the procedure for computing $\text{so}_c : \text{FAC}_c(W) \rightarrow \text{SORT}_c(W)$ considers an initial letter.

Example 6.28. Consider the Coxeter element $c^{\text{ex}} := \tau_1 \tau_2 \tau_3$ of Example 6.2, whose corresponding word is $c^{\text{ex}} w_o(c^{\text{ex}}) = \tau_1 \tau_2 \tau_3 \tau_1 \tau_2 \tau_3 \tau_1 \tau_2 \tau_1$. Let L^{ex} denote the fixed space of $3241 = (13)(34) = (34)(14)$. It is a c^{ex} -noncrossing subspace. To compute $\text{sc}_{c^{\text{ex}}}(L^{\text{ex}})$, we set $s := \psi(\tau_1) = \tau_3$, $c' := s c s = \tau_3 \tau_1 \tau_2$ and $c'' := c s = s c' = \tau_1 \tau_2$. Since L^{ex} is contained in the reflection hyperplane of s , we compute by induction $\text{sc}_{c''}(L^{\text{ex}}) = \{3, 4\}$. Embedding this facet in $c^{\text{ex}} w_o(c^{\text{ex}})$ and adding position 9, we obtain that $\text{sc}_{c^{\text{ex}}}(L^{\text{ex}}) = \{4, 6, 9\}$.

To conclude this section, we summarize in Figure 20 all correspondences obtained in Examples 6.14, 6.16, 6.26, and 6.28.




FIGURE 20. The complete picture of the ongoing example. The upper positions in the facet are drawn in **bold blue**, and below its root configuration is shown.

6.4. Cambrian lattices and fans. Using the connections established in the previous section, we relate in this section the Cambrian lattice with the increasing flip order and the Cambrian fan with the normal fan of the brick polytope.

The c -sortable elements of W form a sublattice and a quotient lattice of the weak order on W , see [Rea06]. This lattice is called *c -Cambrian lattice*. The downward projection used to construct the quotient associates to any element $w \in W$ the maximal c -sortable element below w . It is denoted by $\pi_{\downarrow}^c$, see [RS11, Corollary 6.2]. In [RS11, Theorem 6.3], N. Reading and D. Speyer give a geometric description of the downward projection $\pi_{\downarrow}^c$. This implies first that the bijection so_c^{-1} between c -sortable elements and facets of the subword complex $\mathcal{S}(cw_o(c))$ is the restriction to c -sortable elements of the surjective map κ from W to facets as defined in Section 5.1. It moreover yields the connection between κ and the map $\pi_{\downarrow}^c$.

Theorem 6.29. *Let $v, w \in W$ and let v be c -sortable. Then*

$$\pi_{\downarrow}^c(w) = v \iff \kappa(w) = \kappa(v).$$

Proof. By [RS11, Proposition 5.1, Theorem 6.3] and polarity, $\pi_{\downarrow}^c(w) = v$ if and only if $\mathcal{C}_c(v) \subseteq w(\Phi^+)$. As v is c -sortable, Proposition 6.19 gives us that $\mathcal{C}_c(v) = R(\kappa(v))$. The statement follows with the observation that $R(\kappa(v)) \subseteq w(\Phi^+)$ if and only if $\kappa(w) = \kappa(v)$ by definition. $\square$

Example 6.30. Figure 21 shows the map $\kappa : A_3 \rightarrow \mathcal{S}(\text{cw}_o(c))$, for the Coxeter element $c^{\text{ex}} := \tau_1 \tau_2 \tau_3$ of Example 6.2.

The first consequence of Theorem 6.29 is the connection between the Cambrian lattice and the increasing flip order.

Corollary 6.31. *The bijection so_c is an isomorphism between the increasing flip order on the facets of $\mathcal{S}(\text{cw}_o(c))$ and the Cambrian lattice on c -sortable elements in W .*

Remark 6.32. In fact, Corollary 6.31 was already known for simply-laced types. As pointed out in [CLS11], the structure of the cluster complex as a subword complex can also be deduced from work by K. Igusa and R. Schiffler [IS10, Theorem 2.8]. Together with the work of C. Ingalls and H. Thomas [IT09, Proposition 4.4], one can deduce the previous corollary in simply-laced types.

Remark 6.33. As mentioned above, we conjecture that the increasing flip order on the facets of $\mathcal{S}(Q)$ is a lattice for any realizing word Q . Furthermore, for general words Q , computations in several examples suggest that the increasing flip order still defines such a lattice structure on the facets of the subword complex for which the root configuration is pointed.

The second consequence of Theorem 6.29 is the connection between the Cambrian fan and the normal fan of the brick polytope, which was the heart of the construction of [HLT11].

Corollary 6.34. *The normal fan of the brick polytope $\mathcal{B}(\text{cw}_o(c))$ coincides with the c -Cambrian fan.*

Example 6.35. Figure 22 shows the c^{ex} -Cambrian fan as the normal fan of the brick polytope $\mathcal{B}(c^{\text{ex}}\text{w}_o(c^{\text{ex}}))$, for the Coxeter element $c^{\text{ex}} := \tau_1 \tau_2 \tau_3$ of Example 6.2.

7. CONCLUSION

We proposed in this paper a new approach to the construction of generalized associahedra of finite types, based on the interpretation of cluster complexes in terms of subword complexes [CLS11]. This approach generalizes to all subword complexes of realizing words, thus providing polytopal realizations for a larger class of subword complexes. However, this approach seems to reach its limits for non-realizing words. Indeed, when the word Q is not realizing, the brick polytope does not realize the subword complex $\mathcal{S}(Q)$ (the dimensions do not even match), and may even not be a possible projection of a hypothetical realization of $\mathcal{S}(Q)$ (see [PS12, Proposition 5.9] for an example). Thus, realizations of all spherical subword complexes (or equivalently of all generalized multi-cluster complexes [CLS11]) cannot be achieved using our present approach. For a non-realizing word Q though, the brick polytope $\mathcal{B}(Q)$ still carries relevant information about the flip graph. The root configuration of a facet I still seems to drive the geometry of the cone of the brick polytope at the brick vector of I , and the relevant facets of the subword complex $\mathcal{S}(Q)$ for the brick polytope $\mathcal{B}(Q)$ are thus the facets whose root configuration is pointed. We expect




FIGURE 21. The map κ from $\mathfrak{S}_4$ to the facets of the subword complex $\mathcal{S}(c^{\text{ex}}w_o(c^{\text{ex}}))$ for the Coxeter element $c^{\text{ex}} = \tau_1\tau_2\tau_3$ of Example 6.2. Each node corresponds to a permutation of $\mathfrak{S}_4$ and contains its one-line notation, its c^{ex} -sorting word and its image under κ . The c^{ex} -sortable elements are surrounded by strong boxes. The fibers of κ are represented by shaded regions. The quotient of the weak order by these fibers is the c^{ex} -Cambrian lattice and also the increasing flip order on the facets of $\mathcal{S}(c^{\text{ex}}w_o(c^{\text{ex}}))$. The strong edges between c^{ex} -sortable elements are the edges of the c^{ex} -sorting tree.




FIGURE 22. The normal fan of $\mathcal{B}(c^{\text{ex}}w_o(c^{\text{ex}}))$ is the c^{ex} -Cambrian fan.

that most of the statements concerning the combinatorial and geometric properties of the brick polytope developed in Section 5 can be adapted to brick polytopes of non-realizing words, taking into account only the facets of $\mathcal{S}(Q)$ whose root configuration is pointed. For example, the combinatorics of the brick polytope of any type A spherical subword complex, realizing or not, is described in [PS12].

ACKNOWLEDGEMENTS

The second author wants to thank Pavlo Pylyavskyy for pointing him to [BZ97] which was the starting point for the present paper. We thank Cesar Ceballos, Frédéric Chapoton, Christophe Hohlweg, Jean-Philippe Labbé, Carsten Lange, Nathan Reading, and Hugh Thomas for fruitful discussions on the topic. The first author also thanks Francisco Santos and Michel Pocchiola for their inspiration and collaboration on the premises of brick polytopes in type A . He is also grateful for the opportunity to visit the LaCIM at UQÀM, where this work started. He finally acknowledges the Fields Institute of Toronto for his postdoctoral stay in Canada, and Antoine Deza for his supervision.

REFERENCES

- [BGW97] Alexandre V. Borovik, Israel. M. Gelfand, and Neil White. Coxeter matroid polytopes. *Ann. Comb.*, 1(2):123–134, 1997.
- [BZ97] Arkady Berenstein and Andrei Zelevinsky. Total positivity in Schubert varieties. *Comment. Math. Helv.*, 72(1):128–166, 1997.
- [CFZ02] Frédéric Chapoton, Sergey Fomin, and Andrei Zelevinsky. Polytopal realizations of generalized associahedra. *Canad. Math. Bull.*, 45(4):537–566, 2002. Dedicated to Robert V. Moody.
- [CLS11] Cesar Ceballos, Jean-Philippe Labbé, and Christian Stump. Subword complexes, cluster complexes, and generalized multi-associahedra. Preprint, [arXiv:1108.1776](#), 2011.
- [FZ03] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. II. Finite type classification. *Invent. Math.*, 154(1):63–121, 2003.
- [HL07] Christophe Hohlweg and Carsten E. M. C. Lange. Realizations of the associahedron and cyclohedron. *Discrete Comput. Geom.*, 37(4):517–543, 2007.
- [HLT11] Christophe Hohlweg, Carsten E. M. C. Lange, and Hugh Thomas. Permutahedra and generalized associahedra. *Adv. Math.*, 226(1):608–640, 2011.

- [Hum78] James E. Humphreys. *Introduction to Lie algebras and representation theory*, volume 9 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1978. Second printing, revised.
- [Hum90] James E. Humphreys. *Reflection groups and Coxeter groups*, volume 29 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1990.
- [IS10] Kiyoshi Igusa and Ralf Schiffler. Exceptional sequences and clusters. *J. Algebra*, 323(8):2183–2202, 2010.
- [IT09] Colin Ingalls and Hugh Thomas. Noncrossing partitions and representations of quivers. *Compos. Math.*, 145(6):1533–1562, 2009.
- [KM04] Allen Knutson and Ezra Miller. Subword complexes in Coxeter groups. *Adv. Math.*, 184(1):161–176, 2004.
- [KM05] Allen Knutson and Ezra Miller. Gröbner geometry of Schubert polynomials. *Ann. of Math. (2)*, 161(3):1245–1318, 2005.
- [Pil12] Vincent Pilaud. Greedy flip trees in subword complexes. Preprint, [arXiv:1203.2323](https://arxiv.org/abs/1203.2323), 2012.
- [PP12] Vincent Pilaud and Michel Pocchiola. Multitriangulations, pseudotriangulations and primitive sorting networks. *Discrete Comput. Geom.* (DOI: 10.1007/s00454-012-9408-6), 2012.
- [PS09] Vincent Pilaud and Francisco Santos. Multitriangulations as complexes of star polygons. *Discrete Comput. Geom.*, 41(2):284–317, 2009.
- [PS12] Vincent Pilaud and Francisco Santos. The brick polytope of a sorting network. *European J. Combin.*, 33(4):632–662, 2012.
- [PV96] Michel Pocchiola and Gert Vegter. Topologically sweeping visibility complexes via pseudotriangulations. *Discrete Comput. Geom.*, 16(4):419–453, 1996.
- [Rea04] Nathan Reading. Lattice congruences of the weak order. *Order*, 21(4):315–344 (2005), 2004.
- [Rea06] Nathan Reading. Cambrian lattices. *Adv. Math.*, 205(2):313–353, 2006.
- [Rea07a] Nathan Reading. Clusters, Coxeter-sortable elements and noncrossing partitions. *Trans. Amer. Math. Soc.*, 359(12):5931–5958, 2007.
- [Rea07b] Nathan Reading. Sortable elements and Cambrian lattices. *Algebra Universalis*, 56(3-4):411–437, 2007.
- [RS09] Nathan Reading and David E. Speyer. Cambrian fans. *J. Eur. Math. Soc. (JEMS)*, 11(2):407–447, 2009.
- [RS11] Nathan Reading and David E. Speyer. Sortable elements in infinite Coxeter groups. *Trans. Amer. Math. Soc.*, 363(2):699–761, 2011.
- [RSS03] Günter Rote, Francisco Santos, and Ileana Streinu. Expansive motions and the polytope of pointed pseudo-triangulations. In *Discrete and computational geometry*, volume 25 of *Algorithms Combin.*, pages 699–736. Springer, Berlin, 2003.
- [RSS08] Günter Rote, Francisco Santos, and Ileana Streinu. Pseudo-triangulations — a survey. In *Surveys on discrete and computational geometry*, volume 453 of *Contemp. Math.*, pages 343–410. Amer. Math. Soc., Providence, RI, 2008.
- [SS12] Luis Serrano and Christian Stump. Maximal fillings of moon polyominoes, simplicial complexes, and schubert polynomials. *Electron. J. Combin.*, 19, 2012.
- [Ste11] Salvatore Stella. Polyhedral models for generalized associahedra via coxeter elements. Preprint, [arXiv:1111.1657](https://arxiv.org/abs/1111.1657), 2011.
- [Stu11] Christian Stump. A new perspective on k -triangulations. *J. Combin. Theory Ser. A*, 118(6):1794–1800, 2011.
- [Zie95] Günter M. Ziegler. *Lectures on polytopes*, volume 152 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.

CNRS & LIX, ÉCOLE POLYTECHNIQUE, PALAISEAU
E-mail address: vincent.pilaud@lix.polytechnique.fr
URL: <http://www.lix.polytechnique.fr/~pilaud/>

INSTITUT FÜR ALGEBRA, ZAHLENTHEORIE, DISKRETE MATHEMATIK, UNIVERSITÄT HANNOVER
E-mail address: stump@math.uni-hannover.de
URL: <http://homepage.univie.ac.at/christian.stump/>